

## 32-2017: EDN2 His Recombinant Protein

**Alternative Name :** ET2,PPET2,Endothelin-2,ET-2,Preproendothelin-2,EDN2.

### Description

Source : Escherichia Coli. EDN2 Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 177 amino acids (25-178 a.a.) and having a molecular mass of 19.9kDa. EDN2 is fused to a 23 amino acid His-tag at N-terminus. Endothelin-2 is a Hypoxia-induced Autocrine Survival Factor for Breast Tumor Cells. The synthesis of EDN2 by human kidney carcinoma cells is decreased by EGF. EDN2 is a chemoattractant for macrophages and THP-1 monocytic cells. Chemotaxis towards EDN2 is via the MAPK pathway: p44 and p42 are phosphorylated when THP-1 cells are stimulated with EDN2. Migration to EDN2 is inhibited by hypoxia and by pertussis toxin. EDN2 leads to activation of macrophages. EDN2 shares a similar peptide sequence with chemokines and may signal via a similar receptor and MAPK-mediated pathway. Furthermore, EDN2 expression by tumors may modulate the behavior of macrophages such that activated cells accumulate in areas of hypoxia.

### Product Info

<b>Amount :</b>	20 µg
<b>Purification :</b>	Greater than 85.0% as determined by SDS-PAGE.
<b>Content :</b>	EDN2 protein solution (1mg/ml) containing 20mM Tris-HCl buffer (pH 8.0), 0.4M urea and 10% glycerol.
<b>Storage condition :</b>	Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Avoid multiple freeze-thaw cycles.
<b>Amino Acid :</b>	MGSSHHHHHH SGLVPRGSH MGSQAAATLE QPASSSHAQG THLRLRRCSC SSWLDKECVY FCHLDIIWVN TPEQTAPYGL GNPPRRRRRS LPRRCQCSSA RDPACATFCL RRPWTEAGAV PSRKSPADV QTGKTGATTG ELLQRLDIS TVKSLFAKRQ QEAMREPRST HSRWRKR.

