

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-2063: Procalcitonin Recombinant Protein

Alternative Name: Procalcitonin, PCT.

Description

Source: Escherichia Coli. Procalcitonin Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 116 amino acids and having a molecular mass of 12.8 kDa.The Procalcitonin is purified by standard chromatographic techniques. Procalcitonin is a peptide hormone mainly produced by the C cells of the thyroid and certain endocrine cells of the lung. Under normal expression conditions, procalcitonin is immediately cleaved into three specific fragments, an N terminal residue, calcitonin and katacalcin. Levels of unprocessed procalcitonin rise significantly after bacterial infection. trauma or shock.

Product Info

Amount: $10 \mu g$

Purification: Greater than 97.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

Content: Lyophilized from 10mM Na2PO4 pH-7.5.

Lyophilized procalcitonin although stable at room temperature for 3 weeks, should be stored designated below -18°C. Upon reconstitution procalcitonin should be stored at 4°C between 2-7

Storage condition : desiccated below -18°C. Upon reconstitution procalcitonin should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier

protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Amino Acid: APFRSALESS PADPATLSED EARLLLAALV QDYVQMKASE LEQEQEREGS SLDSPRSKRC GNLSTCMLGT

YTQDFNKFHT FPQTAIGVGA PGKKRDMSSD LERDHRPHVS MPQNAN.

Application Note

It is recommended to reconstitute the lyophilized procalcitonin sterile $18M\tilde{A} \square \hat{A} \odot$ -cm H2O at $100~\tilde{A} \square \hat{A} \mu g/ml$, which can then be further diluted to other aqueous solutions. Human Procalcitonin was tested using Biomerieux reagents on the Min-Vidas. The assay was run at 80 ng/ml with a typical recovery of 80~percent.

