

## 32-2182: BDH2 Recombinant Protein

**Alternative Name :** 3-hydroxybutyrate dehydrogenase type 2,FLJ13261,PRO20933,SDR15C1,U CPA-OR,UNQ6308,dehydrogenase/reductase (SDR family) member 6,Oxidoreductase U CPA,DHRS6,R-beta-hydroxybutyrate dehydrogenase,EFA6R,EC 1.1.1.30.

### Description

Source : Escherichia Coli. BDH2 produced in E.Coli is a single, non-glycosylated polypeptide chain containing 265 amino acids (1-245a.a.) and having a molecular mass of 28.8kDa.BDH2 is fused to a 20 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques. BDH2 is a member of the short-chain dehydrogenases/reductases (SDR) family. BDH2 protein has a significant part in the peripheral utilization of 3-hydroxybutyrate. BDH2 can convert high levels of circulating 3-hydroxybutyrate into acetoacetate due to cytoplasmic localization in high ratio of oxidized NAD+, the NAD+ dependence and the kinetic parameters.

### Product Info

**Amount :** 20 µg  
**Purification :** Greater than 95% as determined by SDS-PAGE.  
**Content :** The BDH2 protein solution (1mg/1ml) is formulated in 20mM Tris-HCl buffer (pH8.0) 0.1M NaCl and 10% glycerol.  
**Storage condition :** Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time.Please avoid freeze thaw cycles.  
**Amino Acid :** MGSSHHHHHH SSSLVPRGSH MGRLDGKVII LTAAQIGIQ AAALAFAREG AKVIATDINE SKLQELEKYP GIQTRVLDVT KKKQIDQFAN EVERLDVLFN VAGFVHHGTV LDCEEKDWDF SMNLNVRSMY LMIKAFLPKM LAQKSGNIIN MSSVASSVKG VVNRCVYSTT KAAVIGLTKS VAADFIQOGI RCNCVCPGTV DTPSLQERIQ ARGNPEEARN DFLKRQKTGR FATAEEIAML CVYLASDESA YVTGNPVIID GGWSL

