

32-2246: Cyclophilin E Recombinant Protein

Alternative Name : Peptidyl-prolyl cis-trans isomerase E, PPIase E, Rotamase E, Cyclophilin-33, PPIE, peptidylprolyl isomerase E, CYP33, Cyclophilin E, CYP-33, MGC3736, MGC111222.

Description

Source : Escherichia Coli. Cyclophilin-E Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 337 amino acids (1-301 a.a.) and having a molecular mass of 37.5 kDa. Cyclophilin-E is fused to 36 amino acids long His Tag at N-terminus and is purified by proprietary chromatographic techniques. Cyclophilin-E is a member of the peptidyl-prolyl cis-trans isomerase (PPIase) family. PPIases catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and speeds up the protein folding. Cyclophilin-E contains a highly conserved cyclophilin domain in addition to a RNA-binding domain. Cyclophilin-E exhibits PPIase activity, protein folding activities and possess RNA-binding activity. Cyclophilin-E contains 2 RNA binding domains at the N-terminal region and a PPIase domain at the C-terminal region.

Product Info

Amount : 25 µg
Purification : Greater than 95.0% as determined by SDS-PAGE.
Content : Cyclophilin-E solution containing 20mM Tris pH-8.
Storage condition : Cyclophilin-E Human Recombinant although stable at 4°C for 1 week, should be stored below -18°C. Please prevent freeze thaw cycles.
Amino Acid : MRGSHHHHHH GMASMTGGQQ MGRDLYDDDD KDRWGSMATT KRVLYVGGLA EEVDDKVLHA
AFIPFGDITD IQIPLDYETE KHRGFAFVEF ELAEDAAAAI DNMNESELF RTIRVNLAKP MRIKEGSSRP
VWSDDDWLKK FSGKTLEENK EEEGSEPPKA ETQEGEPIAK KARSNPQVYM DIKIGNKPAG RIQMLLRSDV
VPMTAENFRC LCTHEKGFGEF KGSSFHRIIP QFMCQGGDFT NHNGTGGKSI YGKKFDDENF ILKHTGPGLL
SMANSGPNTN GSQFFLTCDK TDWLDGKHVV FGEVTEGLDV LRQIEAQGSK DGKPKQKVII ADCGEYV.

Application Note

Specific activity is > 210 nmoles/min/ug, and is defined as the amount of enzyme that cleave 1umole of suc-AAFP-pNA per minute at 1C in Tris-HCl pH8.0 using chymotrypsin.

