

## 32-2249: Cyclophilin H Recombinant Protein

**Alternative Name :** Peptidylprolyl Isomerase H,PPIH,CYPH,CYP20,SnuCyp-20,Peptidyl-prolyl cis-trans isomerase H,PPlase H,Rotamase H,U-snRNP-associated cyclophilin SnuCyp-20,USA-CYP,Small nuclear ribonucleoprotein particle-specific cyclophilin H

### Description

Source : Escherichia Coli. PPIH Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 177 amino acids (1-177) and having a molecular mass of 19.2 kDa. PPIH is purified by proprietary chromatographic techniques. PPIH is a part of the peptidyl-prolyl cis-trans isomerase (PPlase) family. PPlases catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and increase protein folding. PPIH enzyme is a precise factor of the complex that comprises pre-mRNA processing factors PRPF3, PRPF4, and PRPF18, as well as U4/U5/U6 tri-snRNP. PPIH possess PPlase activity and acts as a protein chaperone that mediates the interactions between different proteins inside the spliceosome.

### Product Info

**Amount :** 20 µg  
**Purification :** Greater than 95.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.  
**Content :** 1 mg/ml solution containing 1x PBS pH-7.4 10% glycerol.  
**Storage condition :** PPIH Human Recombinant although stable at 4°C for 1 week, should be stored desiccated below -18°C. Please prevent freeze thaw cycles.  
**Amino Acid :** MAVANSSPVN PVVFFDVSIG GQEVGRMKIE LFADVVPKTA ENFRQFCTGFRKDGVPIGY KGSTFHRVIK DFMIQGGDFV NGDGTGVASI YRGPADENF KLRHSAPGLL SMANSGPSTN GCQFFITCSK CDWLDGKHVV FGKIIDGLLV MRKIENVPTG PNNKPKLPVV ISQCGEM.

### Application Note

Specific activity is > 220 nmoles/min/mg, and is defined as the amount of enzyme that cleaves 1umole of suc-AAFP-pNA per minute at 25C in Tris-Hcl pH8.0 using chymotrypsin.

