

32-2552: MMP 8 Recombinant Protein

Alternative Name : Neutrophil collagenase, EC 3.4.24.34, Matrix metalloproteinase-8, MMP-8, PMNL collagenase, PMNL-CL, HNC, CLG1.

Description

Source : Escherichia Coli. Matrix Metalloproteinase-8 Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain having a molecular mass of 75 kDa. The MMP-8 is purified by proprietary chromatographic techniques. Full-length recombinant human neutrophil pro-collagenase (MMP-8), latent form. Matrix metalloproteinase 8 (MMP-8), or neutrophil collagenase, degrades interstitial collagens, acting preferentially on collagen type I. Increased full-length MMP-8 protein was associated with infiltration into the skin of neutrophils, which are the major cell type that expresses MMP-8. MMP-8 is synthesized and stored in specific granules in neutrophil leukocytes. MMP-8 activity is therefore regulated by factors such as surface-bound ligands (IgG or complement components) that release it through degranulation. Once released and activated through proteolytic or oxidative mechanisms, MMP-8 plays a major role in the connective tissue turnover that accompanies inflammatory processes.

Product Info

Amount : 2.5U
Purification : Greater than 90% as determined by SDS-PAGE.
Content : The MMP-8 protein solution (100 units/ml) in 0.05M Tris-HCl buffer, pH 7.6, 0.2M NaCl, 5mM CaCl₂, 0.0025% NaN₃ and 0.1% BSA.
Storage condition : Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. Avoid multiple freeze-thaw cycles.

Application Note

100 units/ml after activation with APMA by solution assay method. One unit of collagenolytic activity is defined as the cleavage of 1 μ g of collagen per minute by the solution method.

