

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-2781: REV1 Recombinant Protein

Alternative Name:

REV1 Polymerase (DNA Directed), REV1L, Alpha Integrin-Binding Protein 80, Rev1-Like Terminal Deoxycytidyl Transferase, AIBP80, REV1 (Yeast Homolog) - Like, REV1 Homolog (S. Cerevisiae), REV1-Like (Yeast) ,DNA Repair Protein REV1,REV1 Homolog

Description

Source: Escherichia Coli. REV1 Human Recombinant produced in E.coli is a single, non-glycosylated polypeptide chain containing 227 amino acids (51-256) and having a molecular mass of 25.2kDa.REV1 is fused to a 21 amino acid His-tag at Nterminus. REV1 Polymerase (REV1) contains a BRCT domain, which is imperative in protein-protein interactions. REV1 is a deoxycytidyl transferase involved in DNA repair. REV1 assists in the 1st step in the bypass of abasic lesions by the incorporation of a nucleotide opposite the lesion. REV1 is essential for normal generation of mutations by physical and chemical agents. REV1 transfers a dCMP residue from dCTP to the 3'-end of a DNA primer in a template-dependent reaction. A role for the human Rev1-like protein has been suggested as a scaffold which recruits DNA polymerases involved in translesion synthesis (TLS) of damaged DNA.

Product Info

Amount: 20 μg

Purification: Greater than 85.0% as determined by SDS-PAGE.

The REV1 solution (1mg/ml) contains 20mM Tris-HCl buffer (pH 8.0), 0.4M Urea and 10% Content:

glycerol.

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods Storage condition:

of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or

BSA). Avoid multiple freeze-thaw cycles.

Amino Acid: MGSSHHHHHH SSGLVPRGSH MGVAIYVNGY TDPSAEELRK LMMLHGGQYH VYYSRSKTTH IIATNLPNAK

> IKELKGEKVI RPEWIVESIK AGRLLSYIPY QLYTKQSSVQ KGLSFNPVCR PEDPLPGPSN IAKQLNNRVN HIVKKIETEN EVKVNGMNSW NEEDENNDFS FVDLEQTSPG RKQNGIPHPR GSTAIFNGHT PSSNGALKTQ

DCLVPMVNSV ASRLSPA.

