

32-2882: UBE2D1 Recombinant Protein

Alternative Name :

Ubiquitin-conjugating enzyme E2 D1, Stimulator of Fe transport, SFT, UBC4/5 homolog, UbcH5, Ubiquitin carrier protein D1, Ubiquitin-conjugating enzyme E2(17)KB 1, Ubiquitin-conjugating enzyme E2-17 kDa 1, Ubiquitin-protein ligase D1, UBE2D1, SFT, UBC

Description

Source : Escherichia Coli. UBE2D1 produced in E.Coli is a single, non-glycosylated polypeptide chain containing 170 amino acids (1-147 a.a.) and having a molecular mass of 19kDa. UBE2D1 is fused to a 23 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques. UBE2D1 is a member of the ubiquitin-conjugating enzyme family. Ubiquitination involves at least 3 classes of enzymes: ubiquitin-activating enzymes (E1s), ubiquitin-conjugating enzymes (E2s) and ubiquitin-protein ligases (E3s). UBE2D1 is strongly related to a stimulator of iron transport (SFT), and is up-regulated in hereditary hemochromatosis. In addition, UBE2D1 functions in the ubiquitination of the tumor-suppressor protein p53 and the hypoxia-inducible transcription factor HIF1alpha by interacting with the E1 ubiquitin-activating enzyme and the E3 ubiquitin-protein ligases.

Product Info

Amount :	20 µg
Purification :	Greater than 95.0% as determined by SDS-PAGE.
Content :	UBE2D1 solution (1mg/ml) containing 20mM Tris-HCl buffer (pH8.0), 40% glycerol and 0.1M NaCl.
Storage condition :	UBE2D1 Human Recombinant although stable at 4°C for 1 week, should be stored below -18°C. Please prevent freeze thaw cycles.
Amino Acid :	MGSSHHHHHH SGLVPRGSH MGSMALKRIQ KELSDLQRDP PAHCSAGPVG DDLFWQATI MGPPDSAYQG GVFFLTVHFP TDYPFKPPKI AFTTKIYHPN INSNGSICLD ILRSQWSPAL TVSKVLLSIC SLLCDPNPDD PLVPDIAQIY KSDKEKYNRH AREWTQKYAM.

