

32-3217: Bovine Apo Transferrin Native Protein

Alternative Name Serotransferrin,Transferrin,Siderophilin,Beta-1-metal-binding
:
globulin,TF,PRO1557,PRO2086,DKFZp781D0156,Apo Transferrin,ATF.

Description

Source : Bovine Apo Transferrin is a glycoprotein of approximately 77kDa. Transferrin is the iron-transport protein of vertebrate serum and donates iron to cells through interaction with a specific membrane receptor, CD71. Transferrin appears to be indispensable for most cells growing in tissue culture. It is referred to frequently as a growth factor because, in analogy to other growth factor-receptor interactions, proliferating cells express high numbers of transferrin receptors, and the binding of transferrin to their receptors is needed for cells to initiate and maintain their DNA synthesis. Apart from its role as an iron transport protein transferrin acts as a cytokine and has functions that may not be related to its iron-carrying capacity. Bovine Transferrin is a crucial component for the cultivation of mammalian cells in- vitro. Bovine Transferrin is Critical for long-term cells growth in-vitro. Bovine Transferrin is used as detoxificant in media by binding contaminating metal ions. Bovine Transferrin is often used as a nutrient in fermentation media for recombinant protein and biopharmaceutical production. Additional common uses of Bovine Transferrin are Molecular weight, Affinity purification of anti-human transferrin antibodies and also as receptor mediated transfection of molecules such as DNA, into cells.

Product Info

Amount : 150 mg
Purification : Greater than 95.0% as determined by SDS-PAGE.
Content : The protein (1mg/ml) was lyophilized with no additives.
Storage condition : Lyophilized Bovine Apo Transferrin although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Bovine Apo Transferrin should be stored at 4°C between 2-7 days and for future use below -18°C. Please prevent freeze-thaw cycles.

Application Note

It is recommended to reconstitute the lyophilized Apo Transferrin in sterile 18MΩ·cm H₂O not less than 1gr/30ml in 20min. at 20-25C.

