

## 36-1367: Monoclonal Antibody to Cytokeratin 14 (KRT14) (Squamous Cell Marker)(Clone : SPM263)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	SPM263
<b>Application :</b>	IF,IHC
<b>Reactivity :</b>	Human, Mouse, Rat
<b>Gene :</b>	KRT14
<b>Gene ID :</b>	3861
<b>Uniprot ID :</b>	P02533
<b>Format :</b>	Purified
<b>Alternative Name :</b>	KRT14
<b>Isotype :</b>	Mouse IgG3, kappa
<b>Immunogen Information :</b>	A synthetic peptide of 15 amino acid from the C-terminus of human keRatin 14.

### Description

Cytokeratin 14 (CK14) belongs to the type I (or A or acidic) subfamily of low molecular weight keratins and exists in combination with keratin 5 (type II or B or basic). CK14 is found in basal cells of squamous epithelia, some glandular epithelia, myoepithelium, and mesothelial cells. Anti-CK14 is useful in differentiating squamous cell carcinomas from poorly differentiated epithelial tumors. Anti-CK14 is one of the specific basal markers for distinguishing between basal and non-basal subtypes of breast carcinomas. Anti-CK14 is also a good marker for differentiation of intraductal from invasive salivary duct carcinoma by the positive staining of basal cells surrounding the in-situ neoplasm as well as for differentiation of benign prostate from prostate carcinoma. Furthermore, this antibody has been useful in separating oncocytic tumors of the kidney from its renal mimics, and in identifying metaplastic carcinomas of the breast.

### Product Info


<b>Amount :</b>	100 µg
<b>Purification :</b>	Affinity Chromatography
<b>Content :</b>	100 µg in 500 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.
<b>Storage condition :</b>	Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid repeated freeze and thaw cycles.

### Application Note

Immunofluorescence (1-2ug/ml); Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes);



Formalin-fixed, paraffin-embedded human Prostate Carcinoma stained with Cytokeratin 14 Monoclonal Antibody (SPM263).



Formalin-fixed, paraffin-embedded human Prostate Carcinoma stained with Cytokeratin 14 Monoclonal Antibody (SPM263).