

## 32-3449: CD5 Recombinant Protein

**Alternative Name :** CD5 Molecule, CD5, LEU1, CD5 Antigen (P56-62), Lymphocyte Antigen T1/Leu-1, T1, T-Cell Surface Glycoprotein CD5, CD5 Antigen.

### Description

Source : E.coli. CD5 Human Recombinant produced in E.coli is a single, non-glycosylated polypeptide chain containing 371 amino acids (25-372) and having a molecular mass of 41.0 kDa. CD5 is fused to a 23 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques. CD5 is a suitable immunohistochemical marker for T-cells, though not as sensitive as CD3. Approximately 76% of T-cell neoplasms are reported to express CD5, and it is also observed in chronic lymphocytic leukemia, hairy cell leukemia, and mantle cell lymphoma cells. CD5 is commonly lost in cutaneous T-cell lymphoma, and its deficiency can be used as an indicator of malignancy in this condition. CD5's absence in T cell acute lymphoblastic leukemia, though relatively rare, is linked with a poor prognosis. CD5 serves as a receptor in regulating T-cell proliferation.

### Product Info

**Amount :** 20 µg  
**Purification :** Greater than 80% as determined by SDS-PAGE.  
**Content :** The CD5 solution (1mg/ml) contains 20mM Tris-HCl buffer (pH8.0), 10% glycerol and 0.4M Urea.  
**Storage condition :** Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.  
**Amino Acid :** MGSSHHHHHH SSGLVPRGSH MGSRLSWYDP DFQARLTRSN SKCQGQLEVY LKDGWHMVCS QSWGRRSSKQW EDPSQASKVC QRLNCGVPLS LGPFLVITYP QSSIICYGQL GSFSNCSHSR NDMCHSLGLT CLEPQKTTTP TTRPPPTTTP EPTAPRLQL VAQSGGQHCA GVVEFYSGSL GGTISYEAQD KTQDLENFLC>NNLQCGSFLK HLPETEAGRA QDPGEPREHQ PLPIQWKIQN SSCTSLEHCF RKIKPQKSGR VLALLCSGFQ PKVQSRLVGG SSICEGTVEV RQGAQWAALC DSSSARSSLR WEEVCREQQC GSVNSYRVLD AGDPTSRGLF CPHQKLSQCH ELWERNYSYCK KVFVTCQDPN P.

