

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-4016: mlgG His Recombinant Protein

Alternative Name: Ig gamma-2A chain C region, A allele, Immunoglobulin heavy chain gamma polypeptide, Ighq, Igh-1, Igh-1a, 1810060009Rik.

Description

Source: Hi-5 cells. IgG Mouse Recombinant produced in Hi-5 cells is a single polypeptide chain containing 242 amino acids (98-330 a.a) and having a molecular mass of 27.5kDa.lgG is fused to a 6 amino acid His-tag at C-terminus & purified by proprietary chromatographic techniques. Immunoglobulin G (IgG) are antibody molecules. Each IgG is composed of four peptide chains - 2 heavy chains g and 2 light chains. Also, each IgG has 2 antigen binding sites. IgG antibodies are involved in primarily the secondary immune response. The presence of specific IgG, generally, relates to maturation of the antibody response. IgG also has an imperative role in Antibody-dependent cell-mediated cytotoxicity (ADCC) and Intracellular antibody-mediated proteolysis, in which it binds to TRIM21 (the receptor with greatest affinity to IgG in humans) in order to direct marked virions to the proteasome in the cytosol.

Product Info

Amount : 20 μg

Purification : Greater than 90.0% as determined by SDS-PAGE.

Content: IgG protein solution (1mg/ml) containing 20mM Tris-HCl buffer (pH8.0) and 10% glycerol.

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods

Storage condition: of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or

BSA). Avoid multiple freeze-thaw cycles.

Amino Acid: ADPEPRGPTI KPCPPCKCPA PNLLGGPSVF IFPPKIKDVL MISLSPIVTC VVVDVSEDDP DVQISWFVNN

VEVHTAQTQT HREDYNSTLR VVSALPIQHQ DWMSGKEFKC KVNNKDLPAP IERTISKPKG SVRAPQVYVL PPPEEEMTKK QVTLTCMVTD FMPEDIYVEW TNNGKTELNY KNTEPVLDSD GSYFMYSKLR VEKKNWVERN

SYSCSVVHEG LHNHHTTKSF SRTPGKHHHH HH.

