

## 32-4083: Recombinant Human Ku P70/P80

Alternative Name : ATP-dependent DNA helicase 2 subunit 1,ATP-dependent DNA helicase II 70 kDa subunit,Lupus Ku autoantigen protein p70,Ku70,70 kDa subunit of Ku antigen,Thyroid-lupus autoantigen,TLAA,CTC boxbinding factor 75 kDa subunit,CTCBF,CTC75,DNA-repa

## Description

Source : Sf9 insect cells. KuP70/P80 Human Recombinant produced in SF9 insect cells is a glycosylated, polypeptide chain having a molecular mass of 70,638 Dalton for the p70 subunit and 83,528 Dalton for the p80 subunit. Ku (P70/P80) is expressed with a -6xHis tag and purified by proprietary chromatographic techniques. Ku antigen consists of two polypeptides (70 kDa and 80 kDa; the p80 subunit is occasionally called p86). The Ku heterodimer provides DNA-binding and regulatory functions to DNA-dependent protein kinase which is a central enzyme for DNA recombination processes in eucaryotic cells. The Ku protein plays important roles in multiple nuclear processes including DNA repair, V(D)J recombination, telomere maintenance, and the regulation of specific gene transcription. Antibodies to Ku were originally reported in scleroderma-polymyositis overlap syndrome (up to 20% of patients) but have also been detectable in about 20 % of patients with primary pulmonary hypertension (as a complication of overlap), in 5-10 % of SLE and MCTD patients, and in up to 20 % of patients with primary Sjgren's syndrome.In IIF using HEp-2 cells, Ku-positive sera yield a dense, fine granular pattern with nucleolar staining that is typical but not specific for anti-Ku. Detection of Ku antibodies by Western blot may bear false-negative results since the main reactive epitope on the p70/p80 heterodimer is conformation-dependent.

## **Product Info**

Amount :	20 µg
Purification :	Greater than 95% as determined by SDS-PAGE.
Content :	KuP70/P80is supplied in 16mM HEPES buffer pH-8, 160mM NaCl, and 20% glycerol.
Storage condition :	Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. Avoid multiple freeze-thaw cycles.

