w abeomics

32-4337: Recombinant Human Nephrosis 2 Idiopathic Steroid-Resistant

Alternative Name : Nephrosis 2, Idiopathic, Steroid-Resistant (Podocin), PDCN, SRN1, Podocin, NPHS2.

Description

Source : Escherichia Coli. NPHS2 Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain (Val125-Leu383) containing 269 amino acids including a 10 aa His tag at N-terminus. The total calculated molecular mass is 30.2kDa. Nephrosis 2 Idiopathic Steroid-Resistant (NPHS2) is a protein which has a role in the regulation of glomerular permeability, serving as a linker between the plasma membrane and the cytoskeleton. NPHS2 gene mutations cause steroid-resistant nephrotic syndrome.

Product Info

Amount : Purification :	10 μg Greater than 95.0% as determined by SDS-PAGE.
Content :	NPHS2 was filtered (0.4 μm) and lyophilized in 30mM acetate buffer and 1mM dithiothreitol, pH 4.
Storage condition :	Store lyophilized protein at -20°C. Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles. Reconstituted protein can be stored at 4°C for a limited period of time; it does not show any change after two weeks at 4°C.
Amino Acid :	MKHHHHHHASVKVVQEYERV IIFRLGHLLP GRAKGPGLFF FLPCLDTYHK VDLRLQTLEI PFHEIVTKDM FIMEIDAICY YRMENASLLL SSLAHVSKAV QFLVQTTMKR LLAHRSLTEI LLERKSIAQD AKVALDSVTC IWGIKVERIE IKDVRLPAGL QHSLAVEAEA QRQAKVRMIA AEAEKAASES LRMAAEILSG TPAAVQLRYL HTLQSLSTEK PSTVVLPLPF DLLNCLSSPS NRTQGSLPFP SPSKPVEPLN PKKKDSPML.

Application Note

It is recommended to add 0.1M acetate buffer pH 4.0 to prepare a working stock solution of approximately 0.5mg/ml and let the lyophilized pellet dissolve completely. For conversion into higher pH value, we recommend intensive dilution by relevant buffer to a concentration of 10μ g/ml. In higher concentrations the solubility of this antigen is limited. NPHS2 is not sterile! Please filter the product by an appropriate sterile filter before using it in the cell culture.

