

32-4396: Recombinant Human Parkinson Disease Protein 2

Alternative Name :

Parkin RBR E3 Ubiquitin Protein Ligase, Parkinson Juvenile Disease Protein 2, Parkinson Protein 2 E3 Ubiquitin Protein Ligase (Parkin), Parkinson Disease (Autosomal Recessive, Juvenile) 2 Parkin, PRKN, AR-JP, PDJ, E3 Ubiquitin Ligase, LPRS2, EC 6.3.2

Description

Source : Escherichia Coli. PARK2 Human Recombinant produced in E.coli is a single, non-glycosylated polypeptide chain containing 485 amino acids (1-465) and having a molecular mass of 53.8 kDa. PARK2 is fused to a 20 amino acid His-tag at N-terminus. PARK2 is an element in a multiprotein E3 ubiquitin ligase complex which mediates the targeting of substrate proteins for proteasomal degradation. PARK2 alterations can result in Parkinson disease and autosomal recessive juvenile Parkinson disease. Alternative splicing creates multiple transcript variants encoding distinct isoforms.

Product Info

Amount :	20 µg
Purification :	Greater than 80% as determined by SDS-PAGE.
Content :	The PARK2 solution (1mg/ml) contains 20mM Tris-HCl buffer (pH 8.0), 0.4M Urea and 10% glycerol.
Storage condition :	Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.
Amino Acid :	MGSSHHHHHH SSGLVPRGSH MIVFVRFNSS HGFPVEVDS DTSIFQLKEVV AKRQGV PADQ LRVIFAGKEL RNDWTVQNCD LDQQSIVHIV QRPWRKGQEM NATGGDDPRN AAGGCEREPEQ SLTRVDLSSS VLPGDSVGLA VILHTDSRKD SPPAGSPAGR SIYNSFYVYC KGPCQRVQPG KLRVQCSTCR QATLTLTQGP SCWDDVLIPN RMSGECQSPH CPGTSAEFF KCGAHPTSDK ETSVALHLIA TNSRNITCIT CTDVRSPLV FQCNSRHVIC LDCFHLYCVT RLNDRQFVHD PQLGYSLPCV AGCPNSLIKE LHHFRILGEE QYNRYQQYGA EECVLQMGV LCPRPGCGAG LLPEPDQRKV TCEGGNGLGC GFAFCRECKE AYHEGEC SAV FEASGTTTQA YRVD ERAAEQ ARWEAASKET IKKTTKPCPR CHVPVEKNGG CMHMKCPQPQ CRLEWCWNCG CEWNRVCMGD HWFDV

