

32-4445: Recombinant Human Ubiquitin Carboxyl-Terminal Hydrolase L1

Alternative Name : Ubiquitin carboxyl-terminal hydrolase isozyme L1,UCH-L1,EC 3.4.19.12,Ubiquitin thioesterase L1,Neuron cytoplasmic protein 9.5,PGP 9.5,UCHL1,PGP9.5,PARK5.

Description

Source : Escherichia Coli. UCHL1 Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 223 amino acids (1-223 a.a.) and having a molecular mass of 24.8 kDa (molecular weight on SDS-PAGE will appear higher). UCHL1 (PGP9.5) belongs to a gene family whose products hydrolyze small C-terminal adducts of ubiquitin to produce the ubiquitin monomer. Protein Gene Product (PGP9.5) is a neuron specific protein, structurally and immunologically distinct from neuron specific enolase. Human UCHL1 and UCHL3 have an extremely complicated knot structure for a protein, with five knot crossings. It is considered that the knot structure may increase a protein's resistance to degradation in the proteasome. The protein, which has a molecular weight of 27 kDa was first defined by high resolution two dimetional PAGE. Standard immunohistochemical techniques have demonstrated the presence of PGP9.5 in neurons and nerve fibers at all levels of the central and peripheral nervous system, in many neurocrine cells, in segments of the renal tubules, in spermatogonia and leydig cells of the testis, in ova and in some cells of both the pregnant and non pregnant corpus luteum. A point mutation (I93M) in UCHL1 is implicated as the cause of Parkinson's disease in one kindred. On the other hand, a polymorphism (S18Y) in UCHL1 has been found to be associated with a reduced risk for Parkinson's disease. Furthermore, UCHL1 is associated with the Alzheimer's disease, and required for normal synaptic and cognitive function.

Product Info

| | |
|----------------------------|--|
| Amount : | 25 µg |
| Purification : | Greater than 95.0% as determined by SDS-PAGE. |
| Content : | The UCHL1 protein solution (1mg/ml) contains 20mM Tris-HCl buffer (pH 8.0) and 2mM EDTA. |
| Storage condition : | Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Avoid multiple freeze-thaw cycles. |
| Amino Acid : | MQLKPMIENP EMLNKVLSRL GVAGQWRFVD VLGLEESLGSVPAPACALL LLFPLTAQHE NFRKKQIEEL KGQEVSPKVY FMKQTIGNSC GTIGLIHAVA NNQDKLGFED GSVLKQFLSE TEKMSPEDRA KCFEKNEAIQ AAHDAVAQEG QCRVDDKVN FHFILFNNVDG HLYELDGRMP FPNVHGASSE DTLKDAAKV CREFTEREQG EVRFSAVALC KAA. |

Application Note

Specific activity: > 300 pmole/min/ug. Measured by the hydrolysis of Ubiquitin-AMC at pH 8.0, at 37C.

