

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-4636: Recombinant Human Renin, HEK

**Alternative Name :** Renin,Angiotensinogenase,EC 3.4.23.15,HNFJ2,Angiotensin-Forming Enzyme,Renin Precursor Renal,EC 3.4.23.

## **Description**

Source: HEK 293. Renin Human Recombinant produced in HEK cells is a single, glycosylated, polypeptide chain (Leu24-Arg406) containing a total of 393 amino acids, having a calculated molecular mass of 43.7kDa and fused to a 10 aa His tag at C-Terminus. Renin is a highly specific endopeptidase which generates angiotensin I from angiotensinogen in the plasma. angiotensin I is an important regulator of blood pressure and electrolyte balance. Renin initiates a cascade of reactions that produce an elevation of blood pressure and increased sodium retention by the kidney.

## **Product Info**

Amount: 10 μg

**Purification:** Greater than 95.0% as determined by SDS-PAGE.

REN was filtered (0.4µm) and lyophilized from 0.5mg/ml solution in phosphate buffered saline Content:

and 5% (w/v) trehalose.

Store lyophilized protein at -20°C. Aliquot the product after reconstitution to avoid repeated Storage condition:

freezing/thawing cycles. Reconstituted protein can be stored at 4°C for a limited period of time;

it does not show any change after two weeks at 4°C.

Amino Acid: LPTDTTTFKR IFLKRMPSIR ESLKERGVDM ARLGPEWSQP MKRLTLGNTT SSVILTNYMD TQYYGEIGIG

TPPQTFKVVF DTGSSNVWVP SSKCSRLYTA CVYHKLFDAS DSSSYKHNGT ELTLRYSTGT VSGFLSQDII TVGGITVTQM FGEVTEMPAL PFMLAEFDGV VGMGFIEQAI GRVTPIFDNI ISQGVLKEDV FSFYYNRDSE NSOSLGGOIV LGGSDPOHYE GNFHYINLIK TGVWOIOMKG VSVGSSTLLC EDGCLALVDT GASYISGSTS SIEKLMEALG AKKRLFDYVV KCNEGPTLPD ISFHLGGKEY TLTSADYVFQ ESYSSKKLCT LAIHAMDIPP

PTGPTWALGA TFIRKFYTEF DRRNNRIGFA LAR HHHHHHH HHH.

## **Application Note**

It is recommended to add 200µl deionized water to prepare a working stock solution of approximately 0.5 mg/ml and let the lyophilized pellet dissolve completely. REN is not sterile! Please filter the product by an appropriate sterile filter before using it in the cell culture.

