

32-4710: Recombinant Human RCD1 Required for Cell Differentiation1

Alternative Name : Cell differentiation protein RCD1 homolog,Rcd-1,RQCD1,RCD1,CNOT9,CT129,RCD1+.

Description

Source : Escherichia Coli. RQCD1 Human Recombinant fused with a 20 amino acid His tag at N-terminus produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 319 amino acids (1-299 a.a.) and having a molecular mass of 35.7kDa. The RQCD1 is purified by proprietary chromatographic techniques. RQCD1 is a novel transcriptional cofactor which mediates retinoic acid-induced cell differentiation. RQCD1 down-regulates MYB- and JUN-dependent transcription and may have a role in cell differentiation. RQCD1 also has a role in retinoic acid-controlled lung development.

Product Info

Amount :	20 µg
Purification :	Greater than 95.0% as determined by SDS-PAGE.
Content :	The RQCD1 solution (1 mg/ml) contains 20mM Tris-HCl buffer (pH8.0), 0.1M NaCl, 1mM DTT and 10% glycerol.
Storage condition :	RQCD1 should be stored desiccated below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.
Amino Acid :	MGSSHHHHHH SSGLVPRGSH MHSALATAAPV PTTLAQVDRE KIYQWINELS SPETRENALL ELSKKRESVP DLAPMLWHSF GTIAALLQEI VNIYPSINPP TLTAHQSNRV CNALALLQCV ASHPETRSFAF LAAHIPLFLY PFLHTVSKTR PFEYLRRLTSL GVIGALVKTD EQEVINFLLT TEIIPCLLRI MESGSSELSKT VATFILQKIL LDDTGLAYIC QTYERFSHVA MILGKMLVQL SKEPSARLLK HVVRCYLRLS DNPRAREALR QCLPDQLKDT TFAQVLKDDT TTKRWLAQLV KNLQEGQVTD PRGIPLPPQ.

