

## 32-5125: Recombinant Human Triggering Receptor Expressed on Myeloid Cells 2

**Alternative Name :** Triggering receptor expressed on myeloid cells 2, Triggering receptor expressed on monocytes 2, TREM-2, TREM2, Trem2a, Trem2b, Trem2c.

### Description

Source : Escherichia Coli. TREM2 Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 181 amino acids (19-161 a.a) and having a molecular mass of 20.4kDa. TREM2 is fused to a 38 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques. TREM2 is a membrane protein that forms a receptor signaling complex with TYROBP. TREM2 may be involved in chronic inflammation by triggering the production of constitutive inflammatory cytokines. Defects in the TREM2 gene are a source of polycystic lipomembranous osteodysplasia with sclerosing leukoencephalopathy (PLOSL).

### Product Info

**Amount :** 10 µg  
**Purification :** Greater than 90.0% as determined by SDS-PAGE.  
**Content :** TREM2 protein solution (0.25mg/ml) contains 20mM Tris-HCl buffer (pH 8.0), 0.15M NaCl, 10% glycerol and 1mM DTT.  
**Storage condition :** Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.  
**Amino Acid :** MRGSHHHHHH GMASMTGGQQ MGRDLYDDDD KDRWGSMMHN TTVFQGVAGQ SLQVSCP YDS MKHWGRRKAW CRQLGEGGPC QRVVSTHNLW LLSFLRRWNG STAITDDTLG GTLTITLRNL QPHDAGLYQC QSLHGSEADT LRKVLVEVLA DPLDHRDAGD LWFPGESESF EDAHVEHSIS R.

