

## 32-5250: Recombinant Human WNT Inhibitory Factor 1

**Alternative Name :** WIF1,WIF-1,Wnt inhibitory factor 1.

### Description

Source : Hi-5 Cells. WIF1 produced in Hi-5 cells is a single, glycosylated polypeptide chain containing 369 amino acids (29-379 a.a.) and having a molecular mass of 40.5 kDa, (48kDa on SDS-PAGE). WIF1 is fused to 16 amino acid His Tag at C-Terminus and purified by proprietary chromatographic techniques. WIF1 binds to wnt proteins and inhibits their activities. WIF1 plays a role in mesoderm segmentation. WNT proteins are extracellular signaling molecules that take part in the control of embryonic development & cancer. WIF1 protein contains a WNT inhibitory factor (WIF) domain and 5 epidermal growth factor (EGF)-like domains. WIF1 takes part in mesoderm segmentation. WIF1 protein is found to be present in fish, amphibia and mammals. WIF1 is a recurrent target in human salivary gland oncogenesis. Downregulation of WIF1 takes part in the development and progression of pleomorphic adenomas. WIF1 is a tumor suppressor, specifically in nonfunctioning pituitary tumors.

### Product Info

<b>Amount :</b>	5 µg
<b>Purification :</b>	Greater than 80.0% as determined by SDS-PAGE.
<b>Content :</b>	The WIF1 protein solution contains 1X PBS pH 7.4, 10% glycerol and 1mM PMSF.
<b>Storage condition :</b>	Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.
<b>Amino Acid :</b>	ADLGPPQEES LYLWIDAHQA RVLIGFEEDI LIVSEGKMAP FTHDFRKAQQ RMPAIPVNIH SMNFTWQAAG QAEYFYEFLS LRSLDKGIMA DPTVNVPLL TVPHKASVVQ VGFPCLGKQD GVAAFEVDVI VMNSEGNTIL KTPQNAIFFK TCQQAECPPG CRNGGFCNER RICECPDGFH GPHCEKALCT PRCMNGGLCV TPGFCICPPG FYGVNCDKAN CSTTCFNGGT CFYPGKCICP PGLEGEQCEI SKCPQPCRNG GKCIGKSKCK CSKGYQGDLG SKPVCEPGCG AHGTCHPEPNK CQCQEGWHGR HCNKRYEASL IHALRPAGA Q LRQHTPSLKK AEERRDPPE NYIWSGRLVP RGSHHHHHH.

