

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-5370: SARS Nucleocapsid Protein

Description

Source: Escherichia coli. The Recombinant SARS-CoV Nucleocapsid Protein is manufactured with N-terminal fusion HisTag. The Recombinant SARS-CoV Nucleocapsid His-Tagged Fusion Proteinis 47.8 kDa containing 422 amino acid residues of the SARS-CoV Nucleocapsid protein and 15 additional amino acid residues - HisTag. SARS Coronavirus is an enveloped virus containing three outer structural proteins, namely the membrane (M), envelope (E), and spike (S) proteins. Spike (S)-glycoprotein of the virus interacts with a cellular receptor and mediates membrane fusion to allow viral entry into susceptible target cells. Accordingly, S-protein plays an important role in virus infection cycle and is the primary target of neutralizing antibodies.

Product Info

Amount : 5 μg / 20 μg

Purification : Greater than 95% as determined by SDS-PAGE.

Content: Sterile filtered and lyophilized from 0.5 mg/ml in 0.05M Acetate buffer pH4.

Store lyophilized protein at -20°C. Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles. Reconstituted protein can be stored at 4°C for a limited period of time;

it does not show any change after two weeks at 4°C.

Amino Acid: MRGSHHHHHH GMASHMSDNG PQSNQRSAPR ITFGGPTDST DNNQNGGRNG ARPKQRRPQG

LPNNTASWFT ALTQHGKEEL RFPRGQGVPI NTNSGPDDQI GYYRRATRRV RGGDGKMKEL SPRWYFYYLG TGPEASLPYG ANKEGIVWVA TEGALNTPKD HIGTRNPNNN AATVLQLPQG TTLPKGFYAE GSRGGSQASS

RSSSRSRGNS RNSTPGSSRG NSPARMASGG GETALALLLL DRLNQLESKV SGKGQQQQGQ TVTKKSAAEA SKKPRQKRTA TKQYNVTQAF GRRGPEQTQG NFGDQDLIRQ GTDYKHWPQI

AQFAPSASAF FGMSRIGMEV TPSGTWLTYH GAIKLDDKDP QFKDNVILLN KHIDAYKTFP PTEPKKDKKK

KTDEAQPLPQ RQKKQPTVTL LPAADMDDFS RQLQNSMSGA SADSTQA.

Application Note

Add 0.2 ml of 0.1M Acetate buffer pH4 and let the lyophilized pellet dissolve completely. For conversion into higher pH value, we recommend intensive dilution by relevant buffer to a concentration of $10\mu g/ml$. In higher concentrations the solubility of this antigen is limited.

