

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

10-7602: Monoclonal Antibody to VWF (Clone: ABM53D2)

Clonality: Monoclonal **Clone Name:** ABM53D2 Application: FACS.WB Reactivity: Human Gene: **VWF** Gene ID: 7450 **Uniprot ID:** P04275 **Purified** Format:

Alternative Name: von Willebrand Factor Isotype: Mouse IgG1, kappa

Immunogen Information: A partial length recombinant protein (a.a 1710-1939) of VWF was used as the immunogen for

this antibody.

Product Info

Amount : $25 \mu g / 100 \mu g$

Purification: Protein G Chromatography

Content: 25 μg in 50 μl/100 μg in 200 μl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium

azide is highly toxic.

Storage condition : Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

WB: 2-4 μg/ml, FACS: 0.5-1 μg/10⁶




Fig-1: Intracellular flow analysis of VWF antibody on HepG2 cells using 0.5 μ g/10^6 cells of antibody (Clone: ABM53D2). Green represents isotype control; red represents anti-VWF antibody. Goat anti-mouse PE conjugate was used as secondary antibody. (Cells were fixed with 4% paraformaldehyde for 10 min and washed with PBS by centrifuging at 1100 for 5 min followed by permeabilization for 20 min and washed again as mentioned above. Then cell were incubated with primary antibody for 45 min. and after washing the cells twice in PBS, incubated with conjugated secondary antibody for 30 min. Data acquisition was done after washing twice with PBS as mentioned above).



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Fig-2: Western blot analysis of VWF. Anti-VWF antibody (Clone : ABM53D2) was used at 2 $\mu g/ml$ on h Lungs lysate.