

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

10-6014: Monoclonal Antibody to RANKL (Clone: ABM12D6)

Clonality: Monoclonal **Clone Name:** ABM12D6 FACS.WB Application: Reactivity: Human Gene: TNFSF11 Gene ID: 8600 **Uniprot ID:** 014788 Format: **Purified**

Tumor necrosis factor ligand superfamily member 11,Osteoclast differentiation

Alternative Name: factor,Osteoprotegerin ligand,Receptor activator of nuclear factor kappa-B ligand,TNF-related

activation-induced cytokine,CD254

Isotype: Mouse IgG1, Kappa

Immunogen Information: Full length recombinant protein of RANKL was used as the immunogen for this antibody.

Product Info

Amount: 25 μg / 100 μg

Purification: Protein G Chromatography

 $25~\mu g$ in $50~\mu l/100~\mu g$ in $200~\mu l$ PBS containing 0.05% BSA and 0.05% sodium azide. Sodium Content:

azide is highly toxic.

Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid Storage condition:

repeated freeze and thaw cycles.

Application Note

WB: 2-4 μg/ml, FACS: 0.5-1 μg/10⁶




Fig-1: Western blot analysis of RANKL. Anti-RANKL antibody (Clone: ABM12D6) was used at 2 µg/ml on hHeart lysate.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Fig-2: Intra cellular flow analysis of RANKL in Jurkat using $0.5 \mu g/10^6$ cells of antibody (Clone: ABM12D6). Green represents isotype control; red represents anti-RANKL antibody. Goat anti-mouse PE conjugate was used as secondary antibody.




Fig-3: Intra cellular flow analysis of RANKL in HepG2 using 0.5 μ g/10^6 cells of antibody (Clone: ABM12D6). Green represents isotype control; red represents anti-RANKL antibody. Goat anti-mouse PE conjugate was used as secondary antibody.