

## 36-1453: Monoclonal Antibody to MUC1 / EMA / CD227 (Epithelial Marker)(Clone : 139H2)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	139H2
<b>Application :</b>	ELISA,FACS,IF,WB,IHC
<b>Reactivity :</b>	Human, Mouse
<b>Gene :</b>	MUC1
<b>Gene ID :</b>	4582
<b>Uniprot ID :</b>	P15941
<b>Format :</b>	Purified
<b>Alternative Name :</b>	MUC1,PUM
<b>Isotype :</b>	Mouse IgG1, kappa
<b>Immunogen Information :</b>	Human milk-fat globule membranes (HMFGM)

### Description


This MAbs reacts with MUC1. The dominant epitope of this MAbs has not yet been determined. MUC1 is a large cell surface mucin glycoprotein expressed by most glandular and ductal epithelial cells and some hematopoietic cell lineages. It is expressed on most secretory epithelium, including mammary gland and some hematopoietic cells. It is expressed abundantly in lactating mammary glands and over expressed abundantly in >90% breast carcinomas and metastases. Transgenic MUC1 has been shown to associate with all four c-erbB receptors and localize with c-erbB1 (EGFR) in lactating glands. The MUC1 gene contains seven exons and produces several different alternatively spliced variants. The major expressed form of MUC1 uses all seven exons and is a type 1 transmembrane protein with a large extracellular tandem repeat domain. The tandem repeat domain is highly O glycosylated and alterations in glycosylation have been shown in epithelial cancer cells. Antibody to EMA is useful as a pan-epithelial marker for detecting early metastatic loci of carcinoma in bone marrow or liver.

### Product Info


<b>Amount :</b>	100 µg
<b>Purification :</b>	Affinity Chromatography
<b>Content :</b>	100 µg in 500 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.
<b>Storage condition :</b>	Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid repeated freeze and thaw cycles.

### Application Note


ELISA (For coating use Ab at 2-5ug/ml, order Ab without BSA0; ,Flow Cytometry (0.25-0.5ug/million cells); Immunofluorescence (1-2ug/ml); Western Blot (1-2ug/ml); ,Immunohistochemistry (Formalin-fixed) (0.5-1ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes),



Formalin-fixed, paraffin-embedded human Lung Cancer stained with EMA Monoclonal Antibody (139H2).



Formalin-fixed, paraffin-embedded human Breast Cancer stained with EMA Monoclonal Antibody (139H2).



Formalin-fixed, paraffin-embedded human Endometrial Carcinoma stained with EMA Monoclonal Antibody (139H2).