

## 35-1066: Polyclonal Antibody to BAD (Phospho-Ser136)

|                                |  |
|--------------------------------|--|
| <b>Clonality :</b>             | Polyclonal   |
| <b>Application :</b>           | WB,IHC   |
| <b>Reactivity :</b>            | Human,Mouse,Rat  |
| <b>Gene :</b>                  | Bad  |
| <b>Gene ID :</b>               | 12015  |
| <b>Uniprot ID :</b>            | Q61337   |
| <b>Format :</b>                | Purified   |
| <b>Alternative Name :</b>      | Bbc2   |
| <b>Isotype :</b>               | Rabbit IgG   |
| <b>Immunogen Information :</b> | Peptide sequence around phosphorylation site of serine136 (S-R-S(p)-A-P) derived from Mouse BAD. |

### Description

The protein encoded by BAD gene is a member of the BCL-2 family. BCL-2 family members are known to be regulators of programmed cell death. This protein positively regulates cell apoptosis by forming heterodimers with BCL-xL and BCL-2, and reversing their death repressor activity. Proapoptotic activity of this protein is regulated through its phosphorylation. Protein kinases AKT and MAP kinase, as well as protein phosphatase calcineurin were found to be involved in the regulation of this protein. Alternative splicing of this gene results in two transcript variants which encode the same isoform. Wang XQ, et al. (2001). J Biol Chem.276 (48): 44504-44511. Lee YI, et al. (2001). J Biol Chem.276 (20): 16969-16977. Maiti D, et al. (2001). J Biol Chem.276 (1): 329-333.

### Product Info

|                            |  |
|----------------------------|--|
| <b>Amount :</b>            | 50 µl / 100 µl   |
| <b>Content :</b>           | Supplied at 1.0mg/mL in phosphate buffered saline (without Mg <sup>2+</sup> and Ca <sup>2+</sup> ), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. |
| <b>Storage condition :</b> | Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles.                                |

### Application Note

Predicted MW: 23kd, Western blotting: 1:500~1:1000, Immunohistochemistry: 1:50~1:100




Figure 1: Western blot analysis of extracts from HT29 cells untreated or treated with UV using BAD(Phospho-Ser136) Antibody 35-1066 .




Figure 2: Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using BAD(Phospho-Ser136) Antibody 35-1066 (left) or the same antibody preincubated with blocking peptide(right).