

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

35-1114: Polyclonal Antibody to BRCA1 (Phospho-Ser1524)

Clonality: Polyclonal
Application: IHC
Reactivity: Human
Gene: BRCA1
Gene ID: 672
Uniprot ID: P38398

Format: Purified
Alternative Name: RNF53
Isotype: Rabbit IgG

Immunogen Information: Peptide sequence around phosphorylation site of serine 1524 (Y-P-S(p)-Q-E) derived from

Human BRCA1.

Description

The BRCA1-BARD1 heterodimer coordinates a diverse range of cellular pathways such as DNA damage repair, ubiquitination and transcriptional regulation to maintain genomic stability. Acts by mediating ubiquitin E3 ligase activity that is required for its tumor suppressor function. Plays a central role in DNA repair by facilitating cellular response to DNA repair. Required for appropriate cell cycle arrests after ionizing irradiation in both the S-phase and the G2 phase of the cell cycle. Involved in transcriptional regulation of P21 in response to DNA damage. Required for FANCD2 targeting to sites of DNA damage. May function as a transcriptional regulator. Inhibits lipid synthesis by binding to inactive phosphorylated ACACA and preventing its dephosphorylation

Product Info

Amount : 50 μl / 100 μl

Content: Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM

NaCl, 0.02% sodium azide and 50% glycerol.

Storage condition : Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

Predicted MW: 220kd, Immunohistochemistry: 1:50~1:100




Figure 1: Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using BRCA1(Phospho-Ser1524) Antibody 35-1114 (left) or the same antibody preincubated with blocking peptide(right).