

35-1152: Polyclonal Antibody to Zap-70 (Phospho-Tyr319)

Clonality :	Polyclonal
Application :	WB,IHC,IF
Reactivity :	Human,Mouse,Rat
Gene :	ZAP70
Gene ID :	7535
Uniprot ID :	P43403
Format :	Purified
Alternative Name :	SRK,STD,TZK,STCD,ZAP-70
Isotype :	Rabbit IgG
Immunogen Information :	Peptide sequence around phosphorylation site of tyrosine 319 (S-P-Y(p)-S-D) derived from Human Zap-70.

Description

Plays a role in T-cell development and lymphocyte activation. Essential for TCR-mediated IL-2 production. Isoform 1 induces TCR-mediated signal transduction, isoform 2 does not. Salomon AR, et al. (2003) Proc Natl Acad Sci U S A; 100(2): 443-448 Ku GM, et al. (2001) EMBO J; 20(3): 457-465 Tang J, et al. (1999) Proc Natl Acad Sci U S A; 96(17): 9775-9780 Zhao Q, et al. (1996) Mol Cell Biol; 16(12): 6765-6774 Williams BL, et al. (1999) EMBO J; 18(7): 1832-1844 Adjali O, et al. (2005) J Clin Invest; 115(8): 2287-2295.

Product Info

Amount :	50 µl / 100 µl
Content :	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage condition :	Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles.

Application Note

Predicted MW: 70kd, Western blotting: 1:500~1:1000, Immunofluorescence: 1:100~1:200




Figure 1: Western blot analysis of extracts from Jurkat cells using Zap-70(Phospho-Tyr319) Antibody 35-1152 (Lane 2) and the same antibody preincubated with blocking peptide(Lane1).




Figure 2: Immunofluorescence staining of methanol-fixed HeLa cells using Zap-70(Phospho-Tyr319) Antibody 35-1152 .




Figure 3: Western blot analysis of extracts from HL60 cells, treated with calf intestinal phosphatase (CIP), using Zap-70 (Phospho-Tyr319) Antibody 35-1152 .