

## 35-1200: Polyclonal Antibody to VASP (Phospho-Ser157)

|                                |  |
|--------------------------------|--|
| <b>Clonality :</b>             | Polyclonal   |
| <b>Application :</b>           | WB,IHC   |
| <b>Reactivity :</b>            | Human,Rat  |
| <b>Gene :</b>                  | VASP   |
| <b>Gene ID :</b>               | 7408   |
| <b>Uniprot ID :</b>            | P50552   |
| <b>Format :</b>                | Purified   |
| <b>Alternative Name :</b>      | VASP   |
| <b>Isotype :</b>               | Rabbit IgG   |
| <b>Immunogen Information :</b> | Peptide sequence around phosphorylation site of serine 157 (R-V-S(p)-N-A) derived from Human VASP. |

### Description

Ena/VASP proteins are actin-associated proteins involved in a range of processes dependent on cytoskeleton remodeling and cell polarity such as axon guidance, lamellipodial and filopodial dynamics, platelet activation and cell migration. VASP promotes actin filament elongation. It protects the barbed end of growing actin filaments against capping and increases the rate of actin polymerization in the presence of capping protein. VASP stimulates actin filament elongation by promoting the transfer of profilin-bound actin monomers onto the barbed end of growing actin filaments. Plays a role in actin-based mobility of *Listeria monocytogenes* in host cells. Regulates actin dynamics in platelets and plays an important role in regulating platelet aggregation.

### Product Info

|                            |  |
|----------------------------|--|
| <b>Amount :</b>            | 50 $\mu$ l / 100 $\mu$ l   |
| <b>Content :</b>           | Supplied at 1.0mg/mL in phosphate buffered saline (without Mg <sup>2+</sup> and Ca <sup>2+</sup> ), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. |
| <b>Storage condition :</b> | Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles.                                |

### Application Note

Predicted MW: 50kd, Western blotting: 1:500~1:1000, Immunohistochemistry: 1:50~1:100




Figure 1: Western blot analysis of extracts from C6 cells untreated or treated with PMA using VASP(Phospho-Ser157) Antibody 35-1200 .




Figure 2: Immunohistochemical analysis of paraffin-embedded human tonsil carcinoma tissue using VASP(Phospho-Ser157) Antibody 35-1200 (left) or the same antibody preincubated with blocking peptide(right).