

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

35-1227: Polyclonal Antibody to p44/42 MAP Kinase (Phospho-Thr202)

Clonality : Polyclonal **Application :** WB,IHC,IF

Reactivity: Human, Mouse, Rat

Gene: MAPK3
Gene ID: 5595
Uniprot ID: P27361
Format: Purified

Alternative Name: ERK, ERT2, Extracellular signal- regulated kinase 1, Insulin-stimulated MAP2 kinase, MAP

kinase 1

Isotype: Rabbit IgG

Immunogen Information: Peptide sequence around phosphorylation site of threonine 202 (F-L-T(p)-E-Y) derived from

Human p44/42 MAP Kinase.

Description

Involved in both the initiation and regulation of meiosis, mitosis, and postmitotic functions in differentiated cells by phosphorylating a number of transcription factors such as ELK-1. Phosphorylates EIF4EBP1; required for initiation of translation. Phosphorylates microtubule-associated protein 2 (MAP2). Phosphorylates SPZ1 TETE HANNKEN, et al. (2000) Am Soc Nephrol 11:1387-1397 Omar D. PerezNature et al. (2002) Biotechnology 20: 155 - 162 Jingui Yu, et al. (2005) Anesth Analg 101: 315-321 Hironobu Ihn et al.(2000) Immunology 165: 2149-2155

Product Info

Amount : $50 \mu l / 100 \mu l$

Content: Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM

NaCl, 0.02% sodium azide and 50% glycerol.

Storage condition : Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

Predicted MW: 42 44 kd, Western blotting: $1:500\sim1:1000$, Immunohistochemistry: $1:50\sim1:100$, Immunofluorescence: $1:100\sim1:200$




Figure 1: Western blot analysis of extracts from 293 cells untreated or treated with PMA for the indicated times, using p44/42 MAP Kinase(Phospho-Thr202) Antibody 35-1227.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2: Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using p44/42 MAP Kinase (Phospho-Thr202) Antibody 35-1227 (left) or the same antibody preincubated with blocking peptide 51245 (right).




Figure 3: Immunofluorescence staining of methanol-fixed Hela cells showing cytoplasmic, nuclear staining using p44/42 MAP Kinase (Phospho-Thr202) Antibody 35-1227.