

35-1258: Polyclonal Antibody to Integrin beta3 (Phospho-Tyr785)

Clonality :	Polyclonal
Application :	WB
Reactivity :	Human,Mouse,Rat
Gene :	ITGB3
Gene ID :	3690
Uniprot ID :	P05106
Format :	Purified
Alternative Name :	CD61 antigen, GP3A, GPIIIa, ITB3, Platelet membrane glycoprotein IIIa
Isotype :	Rabbit IgG
Immunogen Information :	Peptide sequence around phosphorylation site of tyrosine 785 (I-T-Y(p)-R-G) derived from Human Integrin b3.

Description

Integrin α -V/ β -3 is a receptor for cytotactin, fibronectin, laminin, matrix metalloproteinase-2, osteopontin, osteomodulin, prothrombin, thrombospondin, vitronectin and von Willebrand factor. Integrin α -IIb/ β -3 is a receptor for fibronectin, fibrinogen, plasminogen, prothrombin, thrombospondin and vitronectin. Integrins α -IIb/ β -3 and α -V/ β -3 recognize the sequence R-G-D in a wide array of ligands. Integrin α -IIb/ β -3 recognizes the sequence H-H-L-G-G-A-K-Q-A-G-D-V in fibrinogen gamma chain. Following activation integrin α -IIb/ β -3 brings about platelet/platelet interaction through binding of soluble fibrinogen. This step leads to rapid platelet aggregation which physically plugs ruptured endothelial surface. In case of HIV-1 infection, the interaction with extracellular viral Tat protein seems to enhance angiogenesis in Kaposi's sarcoma lesions.

Product Info

Amount :	50 μ l / 100 μ l
Content :	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage condition :	Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles.

Application Note

Predicted MW: 110kd, Western blotting: 1:500~1:1000




Figure 1: Western blot analysis of extracts from HUVEC cells using Integrin β 3(Phospho-Tyr785) Antibody 35-1258 and the same antibody preincubated with blocking peptide .