

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 35-1284: Polyclonal Antibody to Shc1 (Phospho-Tyr349)

Clonality: Polyclonal Application: WB,IF

**Reactivity:** Human, Mouse

Gene: SHC1
Gene ID: 6464
Uniprot ID: P29353
Format: Purified

Alternative Name: SH2 domain protein C1, SHC, SHC-transforming protein 1, SHCA, Src homology 2 domain-

containing-transforming protein C1

**Isotype:** Rabbit IgG

Immunogen Information: Peptide sequence around phosphorylation site of tyrosine 349 (H-Q-Y(p)-Y-N) derived from

Human Shc1.

## **Description**

Signaling adapter that couples activated growth factor receptors to signaling pathway. Isoform p46Shc and isoform p52Shc, once phosphorylated, couple activated receptor tyrosine kinases to Ras via the recruitment of the GRB2/SOS complex and are implicated in the cytoplasmic propagation of mitogenic signals. Isoform p46Shc and isoform p52Shc may thus function as initiators of the Ras signaling cascade in various non-neuronal systems. Isoform p66Shc does not mediate Ras activation, but is involved in signal transduction pathways that regulate the cellular response to oxidative stress and life span. Isoform p66Shc acts as a downstream target of the tumor suppressor p53 and is indispensable for the ability of stress-activated p53 to induce elevation of intracellular oxidants, cytochrome c release and apoptosis. The expression of isoform p66Shc has been correlated with life span

## **Product Info**

**Amount :** 50 μl / 100 μl

Content: Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM

NaCl, 0.02% sodium azide and 50% glycerol.

**Storage condition :** Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

## **Application Note**

Predicted MW: 46 52 kd, Western blotting: 1:500~1:1000, Immunofluorescence: 1:100~1:200




Figure 1: Western blot analysis of extracts from HepG2 cells untreated or treated with EGF using Shc1(Phospho-Tyr349) Antibody 35-1284.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2: Immunofluorescence staining of methanol-fixed Hela cells using Shc1(Phospho-Tyr349) Antibody 35-1284 .