

35-1303: Polyclonal Antibody to JNK1/JNK2/JNK3 (phospho-Thr183/Tyr185)

Clonality :	Polyclonal
Application :	WB,IF
Reactivity :	Human,Mouse,Rat
Gene :	MAPK8
Gene ID :	5599
Uniprot ID :	P45983/ P45984 P
Format :	Purified
Alternative Name :	Stress-activated protein kinase JNK1, c-Jun N-terminal kinase 1, JNK-46
Isotype :	Rabbit IgG
Immunogen Information :	Peptide sequence around phosphorylation site of Thr183/Tyr185 (M-M-T(p)-P-Y(p)- V - V) derived from Human JNK1/JNK2/JNK3.

Description

Responds to activation by environmental stress and pro-inflammatory cytokines by phosphorylating a number of transcription factors, primarily components of AP-1 such as JUN, JDP2 and ATF2 and thus regulates AP-1 transcriptional activity. In T-cells, JNK1 and JNK2 are required for polarized differentiation of T-helper cells into Th1 cells. By similarity, Phosphorylates heat shock factor protein 4 (HSF4). /Responds to activation by environmental stress and pro-inflammatory cytokines by phosphorylating a number of transcription factors, primarily components of AP-1 such as c-Jun and ATF2 and thus regulates AP-1 transcriptional activity. In T-cells, JNK1 and JNK2 are required for polarized differentiation of T-helper cells into Th1 cells. JNK2 isoforms display different binding patterns: α -1 and α -2 preferentially bind to c-Jun, whereas β -1 and β -2 bind to ATF2. However, there is no correlation between binding and phosphorylation, which is achieved at about the same efficiency by all isoforms. JUNB is not a substrate for JNK2 α -2, and JUND binds only weakly to it. /Responds to activation by environmental stress and pro-inflammatory cytokines by phosphorylating a number of transcription factors, primarily components of AP-1 such as c-Jun and ATF2 and thus regulates AP-1 transcriptional activity. Required for stress-induced neuronal apoptosis and the pathogenesis of glutamate excitotoxicity Davis, R.J. (1999) Biochem Soc Symp 64, 1-12. Ichijo, H. (1999) Oncogene 18, 6087-93. Kyriakis, J.M. and Avruch, J. (2001) Physiol Rev 81, 807-69.

Product Info

Amount :	50 μ l / 100 μ l
Content :	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage condition :	Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles.

Application Note

Predicted MW: 46 54 kd, Western blotting: 1:500~1:1000, Immunofluorescence: 1:100~1:200




Figure 1: Western blot analysis of extracts from C6 cells untreated or treated with anisomycin using JNK1/JNK2/JNK3(phospho-Thr183/Tyr185) Antibody 35-1303 .




Figure 2: Immunofluorescence staining of methanol-fixed HeLa cells using JNK1/JNK2/JNK3(phospho-Thr183/Tyr185) Antibody 35-1303 .




Figure 3: Western blot analysis of extracts from 293 cells, treated with Anisomycin or calf intestinal phosphatase (CIP), using JNK1/JNK2/JNK3 (phospho-Thr183/Tyr185) Antibody 35-1303 .