

## 35-1500: Polyclonal Antibody to Myosin Light Chain 2 (Ab-19)

<b>Clonality :</b>	Polyclonal
<b>Application :</b>	WB,IF
<b>Reactivity :</b>	Human,Mouse,Rat
<b>Gene :</b>	MYL9
<b>Gene ID :</b>	10398
<b>Uniprot ID :</b>	P24844
<b>Format :</b>	Purified
<b>Alternative Name :</b>	LC20,MLC2,MRLC1,MYRL2,MLC-2C
<b>Isotype :</b>	Rabbit IgG
<b>Immunogen Information :</b>	Peptide sequence around aa.17~21 (A-T-S-N-V) derived from Human Myosin Light Chain 2.

### Description

Myosin regulatory subunit that plays an important role in regulation of both smooth muscle and nonmuscle cell contractile activity via its phosphorylation. Implicated in cytokinesis, receptor capping, and cell locomotion Janiak A, et al. (2006) Mol Biol Cell. Apr; 17(4): 1606-1619. Croft DR, et al. (2006) Mol Cell Biol. 2 Jun; 26(12): 4612-4627 Li Z, et al. (2006) Mol Cell Biol. Jun; 26(11): 4240-4256

### Product Info

<b>Amount :</b>	50 $\mu$ l / 100 $\mu$ l
<b>Content :</b>	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg <sup>2+</sup> and Ca <sup>2+</sup> ), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
<b>Storage condition :</b>	Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles.

### Application Note

Predicted MW: 18kd, Western blotting: 1:500~1:1000, Immunofluorescence: 1:100~1:200




Figure 1: Western blot analysis of extracts from 293 and 3T3 cells using Myosin Light Chain 2 (Ab-19) Antibody 35-1500 .




Figure 2: Immunofluorescence staining of methanol-fixed HeLa cells using Myosin Light Chain 2 (Ab-19) Antibody 35-1500 .