

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 35-1515: Polyclonal Antibody to Zap-70 (Ab-319)

Clonality: Polyclonal Application: WB,IHC

Reactivity: Human, Mouse, Rat

Gene: ZAP70 Gene ID: 7535 **Uniprot ID:** P43403 Format: Purified

**Alternative Name:** 70 kDa zeta-associated protein, SRK, Syk-related tyrosine kinase, ZA70, ZAP-70

Rabbit IgG Isotype:

Immunogen Information: Peptide sequence around aa.317~321 (S-P-Y-S-D) derived from Human ZAP70.

## **Description**

ZAP70 is a 70-kD tyrosine phosphoprotein that associates with the zeta chain and undergoes tyrosine phosphorylation following TCR stimulation. The ZAP70 gene is expressed in T- and natural KILLER cells. Protein-Tyrosine Kinases (PTKs) play an integral role in T-cell activation. Stimulation of the T-cell antigen receptor results in tyrosine phosphorylation of a number of cellular substrates. One of these is the TCR-zeta chain, which can mediate the transduction of extracellular stimuli into cellular effector functions Salomon AR, et al. (2003) Proc Natl Acad Sci U S A; 100(2): 443-448 Ku GM, et al. (2001) EMBO J; 20(3): 457-465 Tang J, et al. (1999) Proc Natl Acad Sci U S A; 96(17): 9775-9780 Zhao Q, et al. (1996) Mol Cell Biol; 16(12): 6765-6774 Williams BL, et al. (1999) EMBO J; 18(7): 1832-1844 Adjali O, et al. (2005) J Clin Invest; 115(8): 2287-2295.

## **Product Info**

Amount:  $50 \mu l / 100 \mu l$ 

Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM Content:

NaCl, 0.02% sodium azide and 50% glycerol.

Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid Storage condition:

repeated freeze and thaw cycles.

## **Application Note**

Predicted MW: 70kd, Western blotting: 1:500~1:1000, Immunohistochemistry: 1:50~1:100




Figure 1: Western blot analysis of extracts from HL60 cells using Zap-70(Ab-319) Antibody 35-1515.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2: Immunohistochemical analysis of paraffin-embedded human tonsil tissue using Zap-70(Ab-319) Antibody 35-1515 (left) or the same antibody preincubated with blocking peptide(right).