

36-1463: Monoclonal Antibody to MUC5AC (Mucin 5AC / Gastric Mucin)(Clone : SPM297)

Clonality :	Monoclonal
Clone Name :	SPM297
Application :	FACS,IF,IHC
Reactivity :	Human, Mouse, Rat
Gene :	MUC5AC
Gene ID :	4586
Uniprot ID :	P98088
Format :	Purified
Alternative Name :	MUC5AC,MUC5
Isotype :	Mouse IgG1, kappa
Immunogen Information :	M1 mucin preparation from the fluid of an ovarian mucinous cyst belonging to an O Le(a-b) patient

Description


This MAb recognizes the peptide core of gastric mucin M1 (>1,000kDa) (recently identified as Mucin 5AC). Its epitope is destroyed by beta-mercaptoethanol and proteases but not by periodate treatment. Antibody to gastric mucin M1 reacts with the gastric epithelium of normal human gastrointestinal tract as well as with the precancerous and cancerous colon but not with normal adult colon. It also reacts with fetal colonic mucosa. Resurgence of gastric mucin reactivity during colonic carcinogenesis is due to re-expression of the peptide core of gastric (or fetal colonic) mucins.

Product Info

Amount :	100 µg
Purification :	Affinity Chromatography
Content :	100 µg in 500 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.
Storage condition :	Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid repeated freeze and thaw cycles.

Application Note

Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml); Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);



Formalin-fixed, paraffin-embedded human Gastric Carcinoma stained with MUC5AC Monoclonal Antibody (SPM297).