

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

35-1593: Polyclonal Antibody to Histone H2A.X (Ab-139)

Clonality: Polyclonal
Application: WB,IHC,IF
Reactivity: Human
Gene: H2AFX
Gene ID: 3014
Uniprot ID: P16104
Format: Purified

Alternative Name: H2A.X, H2AFX, H2a/x, HIST5-2AX

Isotype: Rabbit IgG

Immunogen Information: Peptide sequence around aa.137~141 (Q-A-S-Q-E) derived from Human Histone H2A.X.

Description

Variant histone H2A which replaces conventional H2A in a subset of nucleosomes. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling. Required for checkpoint-mediated arrest of cell cycle progression in response to low doses of ionizing radiation and for efficient repair of DNA double strand breaks (DSBs) specifically when modified by C-terminal phosphorylation

Product Info

Amount : $50 \mu l / 100 \mu l$

Content: Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM

NaCl, 0.02% sodium azide and 50% glycerol.

Storage condition : Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

Predicted MW: 15kd, Western blotting: $1:500\sim1:1000$, Immunohistochemistry: $1:50\sim1:100$, Immunofluorescence: $1:100\sim1:200$




Figure 1: Western blot analysis of extracts from HepG2 and 293 cells using Histone H2A.X(Ab-139) Antibody 35-1593 .



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2: Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using Histone H2A.X(Ab-139) Antibody 35-1593 (left) or the same antibody preincubated with blocking peptide(right).




Figure 3: Immunofluorescence staining of methanol-fixed Hela cells using Histone H2A.X(Ab-139) Antibody 35-1593 .