

## 35-1612: Polyclonal Antibody to PKR (Ab-451)

<b>Clonality :</b>	Polyclonal
<b>Application :</b>	IHC,IF
<b>Reactivity :</b>	Human
<b>Gene :</b>	EIF2AK2
<b>Gene ID :</b>	5610
<b>Uniprot ID :</b>	P19525
<b>Format :</b>	Purified
<b>Alternative Name :</b>	ADRB2, E2AK2, EIF2AK2, EIF2aK, PRKR
<b>Isotype :</b>	Rabbit IgG
<b>Immunogen Information :</b>	Peptide sequence around aa.449~453 (K-G-T-L-R) derived from Human PKR.

### Description

Following activation by double-stranded RNA in the presence of ATP, the kinase becomes autophosphorylated and can catalyze the phosphorylation of the translation initiation factor EIF2S1, which leads to an inhibition of the initiation of protein synthesis. Double-stranded RNA is generated during the course of a viral infection. Ingrid K. Ruf, et.al. (2005) J. Virol ; 79: 14562 - 14569. Christy M. Hebner,et.al. (2006) J. Gen. Virol ; 87: 3183 - 3193. Rika van Huizen, et.al. (2003) J. Biol. Chem ; 278: 15558 - 15564.

### Product Info

<b>Amount :</b>	50 µl / 100 µl
<b>Content :</b>	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg <sup>2+</sup> and Ca <sup>2+</sup> ), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
<b>Storage condition :</b>	Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles.

### Application Note

Predicted MW: 68kd, Immunohistochemistry: 1:50~1:100, Immunofluorescence: 1:100~1:200




Figure 1: Immunohistochemical analysis of paraffin-embedded human colon carcinoma tissue using PKR(Ab-451) Antibody 35-1612 (left) or the same antibody preincubated with blocking peptide(right).




Figure 2: Immunofluorescence staining of methanol-fixed HeLa cells using PKR(Ab-451) Antibody 35-1612 .