

## 35-1627: Polyclonal Antibody to GSK3 Alpha/beta(Ab-279/216)

<b>Clonality :</b>	Polyclonal
<b>Application :</b>	WB
<b>Reactivity :</b>	Rat,Mouse,Human
<b>Gene :</b>	GSK3A
<b>Gene ID :</b>	2931
<b>Uniprot ID :</b>	P49840/P49841
<b>Format :</b>	Purified
<b>Alternative Name :</b>	Factor A, GSK-3 alpha/beta, kinase GSK3-alpha/beta
<b>Isotype :</b>	Rabbit IgG
<b>Immunogen Information :</b>	Peptide sequence around aa.277~281/214~218 (V-S-Y-I-C) derived from Human GSK3 alpha/beta.

### Description

Participates in the Wnt signaling pathway. Implicated in the hormonal control of several regulatory proteins including glycogen synthase, MYB and the transcription factor JUN. Phosphorylates JUN at sites proximal to its DNA-binding domain, thereby reducing its affinity for DNA. Phosphorylates MUC1 in breast cancer cells, and decreases the interaction of MUC1 with CTNNB1/beta-catenin. Phosphorylates CTNNB1/beta-catenin. Chin PC, et al. Brain Res Mol Brain Res 2005 Jun 13; 137(1-2): 193-201 Takahashi-Yanaga F, et al. Biochem Biophys Res Commun 2004 Apr 02; 316(2): 411-415 Fan G, et al. J Biol Chem 2003 Dec 26; 278(52): 52432-52436 Liao X, et al. Mol Cancer Ther 2003 Nov; 2(11): 1215-1222

### Product Info

<b>Amount :</b>	50 µl / 100 µl
<b>Content :</b>	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg <sup>2+</sup> and Ca <sup>2+</sup> ), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
<b>Storage condition :</b>	Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles.

### Application Note

Predicted MW: 46, 51kd, Western blotting: 1:500~1:1000




Figure 1: Western blot analysis of extracts from HL60, HT29 and 293 cells using GSK3α/β(Ab-279/216) Antibody 35-1627 .