

## 35-1648: Polyclonal Antibody to TrkB (Ab-705)

<b>Clonality :</b>	Polyclonal
<b>Application :</b>	WB,IF
<b>Reactivity :</b>	Rat,Mouse,Human
<b>Gene :</b>	NTRK2
<b>Gene ID :</b>	4915
<b>Uniprot ID :</b>	Q16620
<b>Format :</b>	Purified
<b>Alternative Name :</b>	TRKB, trk-B, GP145-TrkB
<b>Isotype :</b>	Rabbit IgG
<b>Immunogen Information :</b>	Peptide sequence around aa.703~707 (T-D-Y -Y-R) derived from Human TrkB.

### Description

Receptor for brain-derived neurotrophic factor (BDNF), neurotrophin-3 and neurotrophin-4/5 but not nerve growth factor (NGF). Involved in the development and/or maintenance of the nervous system. This is a tyrosine-protein kinase receptor. Known substrates for the TRK receptors are SHC1, PI-3 kinase, and PLC-gamma-1. Woronowicz A, et al. Glycobiology. 2007 Jan;17(1):10-24. Mojsilovic-Petrovic J, et al. J Neurosci. 2006 Sep 6;26(36):9250-63. Lewis MA, et al. Mol Pharmacol. 2006 Apr;69(4):1396-404. Cai D, et al. Physiol Genomics. 2006 Feb 14;24(3):191-7.

### Product Info

<b>Amount :</b>	50 $\mu$ l / 100 $\mu$ l
<b>Content :</b>	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg <sup>2+</sup> and Ca <sup>2+</sup> ), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
<b>Storage condition :</b>	Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles.

### Application Note

Predicted MW: 145kd, Western blotting: 1:500~1:1000, Immunofluorescence: 1:100~1:200




Figure 1: Western blot analysis of extract from C6 cell and using TrkB(Ab-705) Antibody 35-1648 .




Figure 2: Immunofluorescence staining of methanol-fixed NCI-h446 cells showing cytoplasmic staining using TrkB (Ab-705) Antibody 35-1648 .