

## 35-1677: Polyclonal Antibody to Nanog

<b>Clonality :</b>	Polyclonal
<b>Application :</b>	WB
<b>Reactivity :</b>	Human,Mouse
<b>Gene :</b>	NANOG
<b>Gene ID :</b>	79923
<b>Uniprot ID :</b>	Q9H9S0
<b>Format :</b>	Purified
<b>Alternative Name :</b>	Homeobox transcription factor Nanog
<b>Isotype :</b>	Rabbit IgG
<b>Immunogen Information :</b>	Peptide sequence around aa.137~141 (K-Q-V-K-T) derived from Nano g

### Description

Transcription regulator involved in inner cell mass and embryonic stem (ES) cells proliferation and self-renewal. Imposes pluripotency on ES cells and prevents their differentiation towards extraembryonic endoderm and trophectoderm lineages. Blocks bone morphogenetic protein-induced mesoderm differentiation of ES cells by physically interacting with SMAD1 and interfering with the recruitment of coactivators to the active SMAD transcriptional complexes By similarity. Acts as a transcriptional activator or repressor By similarity. Binds optimally to the DNA consensus sequence 5'-TAAT[GT][GT]-3' or 5'-[CG][GA][CG]C[GC]ATTAN[GC]-3' By similarity. When overexpressed, promotes cells to enter into S phase and proliferation

### Product Info

<b>Amount :</b>	50 µl / 100 µl
<b>Content :</b>	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg <sup>2+</sup> and Ca <sup>2+</sup> ), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
<b>Storage condition :</b>	Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles.

### Application Note

Predicted MW: 42kd, Western blotting: 1:1000




Figure 1: Western blot analysis of extracts from human Umbilical cord mesenchymal stem cell using Nano g Antibody 35-1677 .