

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

35-1747: Polyclonal Antibody to FAK (Ab-576/577)

Clonality: Polyclonal Application: WB,IHC

Reactivity: Human, Mouse, Rat

Gene : PTK2
Gene ID : 5747
Uniprot ID : Q05397
Format : Purified

Alternative Name: FADK 1, FAK1, PTK2

Isotype: Rabbit IgG

Immunogen Information: Peptide sequence around aa.574~578/575~579 (T-Y-Y-K-A) derived from Human FAK.

Description

Non-receptor protein-tyrosine kinase implicated in signaling pathways involved in cell motility, proliferation and apoptosis. Activated by tyrosine-phosphorylation in response to either integrin clustering induced by cell adhesion or antibody cross-linking, or via G-protein coupled receptor (GPCR) occupancy by ligands such as bombesin or lysophosphatidic acid, or via LDL receptor occupancy. Plays a potential role in oncogenic transformations resulting in increased kinase activity.

Product Info

Amount : 50 μl / 100 μl

Content: Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM

NaCl, 0.02% sodium azide and 50% glycerol.

Storage condition : Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

Predicted MW: 125kd, Western blotting: 1:500~1:1000, Immunohistochemistry: 1:50~1:100




Figure 1: Western blot analysis of extracts from HT29 and Hela cells using FAK(Ab-576/577) Antibody 35-1747 .



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2: Immunohistochemical analysis of paraffin-embedded human lung carcinoma tissue using FAK(Ab-576/577) Antibody 35-1747 (left) or the same antibody preincubated with blocking peptide(right).