

## 12-1034: Recombinant Rabbit Monoclonal Antibody to CD31 / PECAM-1 (Endothelial Cell Marker)(Clone : C31/1395R)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	C31/1395R
<b>Application :</b>	WB,FACS,IHC
<b>Reactivity :</b>	Human
<b>Gene :</b>	PECAM1
<b>Gene ID :</b>	5175
<b>Uniprot ID :</b>	P16284
<b>Format :</b>	Purified
<b>Alternative Name :</b>	EndoCAM, PECA1, Platelet Endothelial Cell Adhesion Molecule 1, GPIIA'
<b>Isotype :</b>	Rabbit IgG
<b>Immunogen Information :</b>	Human recombinant full-length CD31 protein

### Description

CD31 (PECAM-1) is a transmembrane glycoprotein member of the immunoglobulin supergene family of adhesion molecules. CD31 is expressed by stem cells of the hematopoietic system and is primarily used to identify and concentrate these cells for experimental studies as well as for bone marrow transplantation. Anti-CD31 has shown to be highly specific and sensitive for vascular endothelial cells. Staining of nonvascular tumors (excluding hematopoietic neoplasms) is rare. CD31 MAbs react with normal, benign, and malignant endothelial cells which make up blood vessel lining. The level of CD31 expression can help to determine the degree of tumor angiogenesis, and a high level of CD31 expression may imply a rapidly growing tumor and potentially a predictor of tumor recurrence.

### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Purification :</b>	Purified Ab with BSA and Azide at 200ug/ml
<b>Content :</b>	200ug/ml of Ab purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid repeated freeze and thaw cycles.

### Application Note

Western Blot (1-2 µg/ml); Flow Cytometry (1-2 µg/million cells); Immunohistochemistry (Formalin-fixed) (1-2 µg/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes)




Figure 1: Formalin-fixed, paraffin-embedded human Angiosarcoma stained with CD31 Rabbit Recombinant Monoclonal Antibody (C31/1395R).




Figure 2: SDS-PAGE Analysis Purified CD31 Rabbit Recombinant Monoclonal Antibody (C31/1395R).