

## 12-1052: Recombinant Rabbit Monoclonal Antibody to CD34 (Hematopoietic Stem Cell & Endothelial Marker)(Clone : HPCA1/1806R)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	HPCA1/1806R
<b>Application :</b>	FACS,IHC
<b>Reactivity :</b>	Human, Rat
<b>Gene :</b>	CD34
<b>Gene ID :</b>	947
<b>Uniprot ID :</b>	P28906
<b>Format :</b>	Purified
<b>Alternative Name :</b>	Hematopoietic Progenitor Cell Antigen, HPCA1, Mucosialin
<b>Isotype :</b>	Rabbit IgG
<b>Immunogen Information :</b>	Recombinant full-length human HPCA1 protein

### Description

This antibody recognizes a transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34. Its expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, it is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. CD34 expression is also found in vascular endothelium. Additionally, proliferating endothelial cells overexpress this molecule than the non-proliferating endothelial cells. Anti-CD34 labels > 85% of angiosarcoma and Kaposi s sarcoma, but shows low specificity.

### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Purification :</b>	Purified Ab with BSA and Azide at 200ug/ml
<b>Content :</b>	200ug/ml of recombinant MAb purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid repeated freeze and thaw cycles.

### Application Note

Flow Cytometry (1-2 $\times$ 10<sup>5</sup> µg/million cells); Immunohistochemistry (Formalin-fixed) (2-4 $\times$ 10<sup>5</sup> µg/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes)




Figure 1: Formalin-fixed, paraffin-embedded human Tonsil stained with CD34 Rabbit Recombinant Monoclonal Antibody (HPCA1/1806R).




Figure 2: SDS-PAGE Analysis of Purified CD34 Rabbit Recombinant Monoclonal Antibody (HPCA1/1806R).