

**12-1055: Recombinant Rabbit Monoclonal Antibody (Negative Control for Rabbit Monoclonal Antibodies) (Clone : NCRBM/1520R)**

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	NCRBM/1520R
<b>Application :</b>	ELISA,FACS,IF,IP,WB,IHC
<b>Reactivity :</b>	All species
<b>Format :</b>	Purified
<b>Isotype :</b>	Rabbit IgG, kappa
<b>Immunogen Information :</b>	Recombinant full-length human KRT7 protein

**Description**

To serve as a true negative control for various procedures, this rabbit monoclonal IgG should be used under the identical conditions and at the same dilution as the rabbit monoclonal antibody being used for a positive reaction.

**Product Info**

<b>Amount :</b>	20 µg / 100 µg
<b>Purification :</b>	Purified Ab with BSA and Azide at 200ug/ml
<b>Content :</b>	200ug/ml of Ab purified by Protein A Column. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid repeated freeze and thaw cycles.

**Application Note**

ELISA; Flow Cytometry; Immunofluorescence; Immunoprecipitation; Western Blot (1-2 µg/ml); Immunohistochemistry (Frozen); Immunohistochemistry (Formalin-fixed) (1-2 µg/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes)