

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

#### 90-2125: Human TIMP-2 (Metalloproteinase Inhibitor-2) Pre-Coated ELISA Kit

**Application:** ELISA

## **Description**

This assay employs an antibody specific for Anti-human TIMP-2 coated on a 96-well plate. Standards and samples are pipetted into the wells and TIMP-2 present in a sample is bound to the wells by the immobilized antibody. The wells are washed and biotinylated anti-human TIMP-2 antibody is added. After washing away unbound biotinylated antibody, HRP-conjugated streptavidin is pipetted to the wells. The wells are again washed, a TMB substrate solution is added to the wells and color develops in proportion to the amount of TIMP-2 bound. The Stop Solution changes the color from blue to yellow, and the intensity of the color is measured at 450 nm.

# **Kit Components**

Item	Specifications	Storage
96 well Strip ELISA Plate	8 × 12 well	4°C
Lyophilized Standard	2 vials	-20°C
Sample and Standard Dilution Buffer	20 ml	4°C
Biotinylated Detection Antibody for h TIMP-2	120 μl	4°C (Protected from light)
Antibody Dilution Buffer	10 ml	4°C
HRP Conjugated Streptavidin	120 μl	4°C (Protected from light)
Streptavidin Dilution Buffer	10 ml	4°C
TMB Substrate	10 ml	4°C
Stop Solution	10 ml	4°C
25X Wash Buffer	30 ml	4°C
Plate Sealer	5 pieces	
Product Manual	1	

### **Product Info**

**Amount:** 1 × 96 Tests

**Content :** 1 × 96 well Format (96 tests)



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

**Storage condition :** Please refer to the Manual

## **Application Note**

This immunoassay kit allows for the in vitro quantitative determination of Human TIMP-2 concentrations in serum, plasma and other biological fluids.

Detection Range: 0.156 - 10 ng/ml

Sensitivity: < 0.094 ng/ml




Fig-1: Human TIMP-2 Standard Curve.