

## 34-1019: Monoclonal Antibody to Calbindin(Clone: 5A9)

|                                |   |
|--------------------------------|---|
| <b>Clonality :</b>             | Monoclonal  |
| <b>Clone Name :</b>            | 5A9   |
| <b>Application :</b>           | WB  |
| <b>Reactivity :</b>            | Human   |
| <b>Gene :</b>                  | CALB1   |
| <b>Gene ID :</b>               | 793   |
| <b>Uniprot ID :</b>            | P05937  |
| <b>Format :</b>                | Purified  |
| <b>Alternative Name :</b>      | Calbindin D28,D-28K,Vitamin D-dependent calcium-binding protein, avian-type |
| <b>Isotype :</b>               | Mouse, IgG2a  |
| <b>Immunogen Information :</b> | Full length recombinant human protein                                       |

### Product Info

|                            |   |
|----------------------------|---|
| <b>Amount :</b>            | 50 µl / 100 µl  |
| <b>Content :</b>           | Antibody is supplied as an aliquot of 1 mg/ml of affinity purified antibody   |
| <b>Storage condition :</b> | Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid repeated freeze and thaw cycles. |

### Application Note

WB: 1:5,000. IF/ICC or IHC: 1:5,000.




Figure-1: Immunofluorescent analysis of rat brain cerebellum section stained with mouse mAb to calbindin,(34-1019), dilution 1:2,000, in red, and costained with rabbit pAb to GFAP,(34-1042), dilution 1:5,000 in green. The blue is Hoechst staining of nuclear DNA. Following transcatheterial perfusion with 4% paraformaldehyde, brain was post fixed for 24 hours, cut to 45µM, and free-floating sections were stained with the above antibodies. The (34-1019) calbindin antibody prominently labels the dendrites and perikarya of Purkinje cells in the molecular layer of the cerebellum. The GFAP antibody stains the processes of Bergmann glia in the molecular layer and astroglia in the granular and white matter layers of cerebellum.




Figure-2: Western blot analysis of different neuronal tissue lysates using mouse mAb to calbindin,(34-1019), dilution 1:5,000: [1] protein standard, [2] rat cerebellum, [3] pig hippocampus, and [4] cow cerebellum. Bands at about 26kDa correspond to calbindin protein, heavily expressed in the cerebellum.