

## 34-1054: Polyclonal Antibody to $\alpha$ -internexin/NF66

<b>Clonality :</b>	Polyclonal
<b>Application :</b>	ICC/IF,IHC,WB
<b>Reactivity :</b>	Human, Rat, Mouse, Cow, Pig, Horse
<b>Gene :</b>	INA
<b>Gene ID :</b>	9118
<b>Uniprot ID :</b>	Q16352
<b>Format :</b>	Conc. IgY prep.
<b>Alternative Name :</b>	66 kDa neurofilament protein,Neurofilament 5
<b>Isotype :</b>	Chicken, IgY
<b>Immunogen Information :</b>	Purified recombinant rat Alpha -internexin expressed in and purified from E. coli.

### Product Info

<b>Amount :</b>	50 $\mu$ l / 100 $\mu$ l
<b>Content :</b>	Antibody is supplied as an aliquot of concentrated IgY prep.
<b>Storage condition :</b>	Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid repeated freeze and thaw cycles.

### Application Note

Western blot: 1:10,000. ICC/IF and IHC: 1:1,000.




Figure-1: Immunofluorescent analysis of a rat cerebellum section stained with chicken pAb to  $\alpha$ -internexin, (34-1054), dilution 1:5,000 in red, and costained with mouse mAb to MBP, 34-1072, 1:5,000 in green. The blue is Hoechst staining of nuclear DNA. Following transcardial perfusion of the rat with 4% paraformaldehyde, brain was post fixed for 24 hours, cut to 45 $\mu$ m, and free-floating sections were stained with the above antibodies. The  $\alpha$ -internexin antibody selectively stains axons and dendrites of neuronal cells, in particular Purkinje cells, parallel fibers and the axons of granule cells, while the MBP antibody stains myelin sheaths around axons.




Figure-2: Western blot analysis of different tissue lysates using chicken pAb to  $\alpha$ -Internexin, (34-1054), dilution 1:10,000 in green: [1] protein standard (red), [2] rat brain, [3] rat spinal cord, [4] mouse brain, [5] mouse spinal cord, [6] cow spinal cord and [7] pig spinal cord. 34-1054 antibody reveals the  $\alpha$ -internexin protein with apparent molecular weight of 64 to 66kDa, with some variability among different species.