

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 10-9548: Recombinant Rabbit Monoclonal Antibody to Acetylated Histone H3 Lysine 9 (K9ac) (Clone: RM161)(Discontinued)

Clonality: Monoclonal Clone Name: RM161

**Application:** WB,ELISA,Multiplex,ChIP,ICC,IHC

Reactivity: All Species
Gene: H3F3A
Gene ID: 3020
Uniprot ID: P84243
Format: Purified
Alternative Name: Histone H3.3
Isotype: Rabbit IgG

Immunogen Information: An acetyl-peptide corresponding to the Acetyl-Histone H3 (Lys9)

## **Product Info**

Amount:  $100 \mu g$ 

**Purification:** Protein A affinity purified from an animal origin-free culture supernatant **Content:** 1 mg/ml in 50% Glycerol/PBS with 1% BSA and 0.09% sodium azide

**Storage condition :** Store at -20°C. Avoid repeated freeze and thaw cycles.

## **Application Note**

Clone RM161 reacts to Histone H3 acetylated at Lysine 9 (K9ac). No cross reactivity with other acetylated Lysines in histone H3. Western Blot:  $0.25 \text{ $\tilde{A}$} \square \text{$\tilde{A}$} \mu \text{g/ml}$ ; ICC:  $0.5 \text{ $\tilde{A}$} \square \text{$\tilde{A}$} \mu \text{g/ml}$ ; ChIP:  $2 \text{ $\tilde{A}$} \square \text{$\tilde{A}$} \mu \text{g/ml}$ ; IHC:  $0.1 \text{ $\tilde{A}$} \square \text{$\tilde{A}$} \mu \text{g/ml}$ ;  $2 \text{ $\tilde{A}$} \square \text{$\tilde{A}$} \mu \text{g/ml}$ ; ELISA:  $2 \text{ $\tilde{A}$} \square \text{$\tilde{A}$} \mu \text{g/ml}$ ; Multiplex:  $2 \text{ $\tilde{A}$} \square \text{$\tilde{A}$} \mu \text{g/ml}$ ;  $2 \text{ $\tilde{A}$} \square \text{$\tilde{A}$} \mu \text{g/ml}$ ; Multiplex:  $2 \text{ $\tilde{A}$} \square \text{$\tilde{A}$} \mu \text{g/ml}$ ;  $2 \text{ $\tilde{A}$} \square \text{$\tilde{A}$} \mu \text{g/ml}$ ; Multiplex:  $2 \text{ $\tilde{A}$} \square \text{$\tilde{A}$} \mu \text{g/ml}$ ;  $2 \text{ $\tilde{A}$} \square \text{$\tilde{A}$} \mu \text{g/ml}$ ; Multiplex:  $2 \text{ $\tilde{A}$} \square \text{$\tilde{A}$} \mu \text{g/ml}$ ;  $2 \text{ $\tilde{A}$} \square \text{$\tilde{A}$} \mu \text{g/ml}$ ; Multiplex:  $2 \text{ $\tilde{A}$} \square \text{$\tilde{A}$} \mu \text{g/ml}$ ;  $2 \text{ $\tilde{A}$} \square \text{$\tilde{A}$} \mu \text{g/ml}$ ; Multiplex:  $2 \text{ $\tilde{A}$} \square \text{$\tilde{A}$} \mu \text{g/ml}$ ;  $2 \text{ $\tilde{A}$} \square \text{$\tilde$ 




Figure 1: Clone: RM161 specifically reacts to Histone H3 acetylated at Lysine 9 (K9ac). No cross reactivity with acetyl-ated Lysine 4 (K4ac), Lysine 14 (K14ac), Lysine 18 (K18ac), Lysine 23 (K23ac), Lysine 27 (K27ac), Lysine 36 (K36ac), Lysine 56 (K56ac), Lysine 79 (K79ac), or Lysine 122 (K122) in histone H3.





Figure 2: Western Blot of acid extracts from HeLa cells untreated (-) or treated with sodium butyrate (+), using Clone: RM161 at 0.25 µg/ml, showed a band of histone H3 acetylated at Lysine 9 in treated HeLa.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com



\* abeomics

Figure 3: Sandwich ELISA against acetylated histone H3 at Lys 9 using HeLa whole cell lysate, treated or untreated with Sodium Butyrate. Using anti-H3CT (Clone: RM188, 1  $\mu$ g/ml) as the capture antibody and biotinylated anti-H3K9ac (Clone: RM161,1  $\mu$ g/ml) as the detection antibody.




Figure 4: ChIP performed on HeLa cells using H3K9ac antibody (Clone: RM161, 5  $\mu$ g). Real-time PCR was performed using primers specific to the gene indicated.




Figure 5: Immunocytochemistry of HeLa cells treated with sodium butyrate, using Acetyl-Histone H3 (Lys9) Rabbit mAb Clone: RM161 (red). Actin filaments have been labeled with fluorescein phalloidin (green).




Figure 6: Immunohistochemistry staining of HepG2 cells usinganti-Acetyl-Histone H3 (Lys9) antibody, Clone: RM161.