

## 10-6577: Mouse Monoclonal Antibody to HGF (Clone: 489CT6.12.6)(Discontinued)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	489CT6.12.6
<b>Application :</b>	WB
<b>Reactivity :</b>	Human
<b>Gene :</b>	HGF
<b>Gene ID :</b>	3082
<b>Uniprot ID :</b>	P14210
<b>Format :</b>	Purified
<b>Alternative Name :</b>	Hepatocyte growth factor, Hepatopoietin-A, Scatter factor, SF, Hepatocyte growth factor alpha chain, Hepatocyte growth factor beta chain, HGF, HPTA
<b>Isotype :</b>	Mouse IgG1
<b>Immunogen Information :</b>	Synthetic Peptide

### Description

Hepatocyte growth factor regulates cell growth, cell motility, and morphogenesis by activating a tyrosine kinase signaling cascade after binding to the proto-oncogenic c-Met receptor. Hepatocyte growth factor is secreted by mesenchymal cells and acts as a multi-functional cytokine on cells of mainly epithelial origin. Its ability to stimulate mitogenesis, cell motility, and matrix invasion gives it a central role in angiogenesis, tumorogenesis, and tissue regeneration. It is secreted as a single inactive polypeptide and is cleaved by serine proteases into a 69-kDa alpha-chain and 34-kDa beta-chain. A disulfide bond between the alpha and beta chains produces the active, heterodimeric molecule. The protein belongs to the plasminogen subfamily of S1 peptidases but has no detectable protease activity. Alternative splicing of this gene produces multiple transcript variants encoding different isoforms. [provided by RefSeq].

### Product Info

<b>Amount :</b>	80 µl / 400 µl
<b>Purification :</b>	Protein G Chromatography
<b>Content :</b>	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
<b>Storage condition :</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term store at -20°C in small aliquots to prevent freeze-thaw cycles.

### Application Note

WB~1:1000




Figure 1: All lanes : Anti-Human HGF (10-6577) at 1:1000 dilution with Lane 1: HepG2 whole cell lysate and Lane 2: Ramos whole cell lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 83 kDa.