

10-12502: Mouse Monoclonal Antibody to Bcl2(Clone :BS94)

Clonality :MonoclonalClone Name :BS94Application :IHCReactivity :HumanGene :BCL2Gene ID :596Uniprot ID :P10415

Product Info

Amount :	0.1 ml / 0.5 ml
Content :	TRIS with 0.03% sodium azide, pH7.2
Storage condition :	Store at 4°C

Application Note

Immunohistochemical Analysis :-1:200




Figure-1: Tonsil (Figure-1) and lymph node sections (Figure-2, figure-3) have been stained using BCL2 antibody (Clone: BS94) with 1:200 dilution. All peripheral lymphocytes should be labelled and the germinal center should be negatively stained from tonsil sections. Only the few scattered T-cells should be stained (Figure-2, figure-3). Mantle cell lymphoma stained strongly with membranous staining pattern.

Figure-2: Tonsil (Figure-1) and lymph node sections (Figure-2, figure-3) have been stained using BCL2 antibody (Clone: BS94) with 1:200 dilution. All peripheral lymphocytes should be labelled and the germinal center should be negatively stained from tonsil sections. Only the few scattered T-cells should be stained (Figure-2, figure-3). Mantle cell lymphoma stained strongly with membranous staining pattern.

w abeomics

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com



Figure-3: Tonsil (Figure-1) and lymph node sections (Figure-2, figure-3) have been stained using BCL2 antibody (Clone: BS94) with 1:200 dilution. All peripheral lymphocytes should be labelled and the germinal center should be negatively stained from tonsil sections. Only the few scattered T-cells should be stained (Figure-2, figure-3). Mantle cell lymphoma stained strongly with membranous staining pattern.